Keio Associated Repository of Academic resouces

Relational Action of the Actio	
Title	Antihypertensive and diuretic effects of YM-09730-5, a new calcium antagonist, in stroke-prone spontaneously hypertensive rats
Sub Title	
Author	川島, 紘一郎(Kawashima, Koichiro) 戸田, 浜朗(Toda, Hamao) 大畑, 尚代(Ohata, Hisayo) 藤本, 和子(Fujimoto, Kazuko) 鈴木, 岳之(Suzuki, Takeshi) 稲垣, 治(Inagaki, Osamu)
Publisher	共立薬科大学
Publication year	1991
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.36 (1991.),p.48-48
JaLC DOI	
Abstract	
Notes	<b>沙</b> 録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-0000036-0048

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

## Antihypertensive and Diuretic Effects of YM-09730-5, a new Calcium Antagonist, in Stroke-Prone Spontaneously Hypertensive Rats\*

Koichiro Kawashima, Hamao Toda, Hisayo Oohata, Kazuko Fujimoto, Takeshi Suzuki and Osamu Inagaki\*\*

川島紘一郎, 戸田浜朗, 大畑尚代, 藤本和子, 鈴木岳之, 稲垣 治\*\*,

- 1. Effects of consecutive administration of YM-09730-5, (3S)-1-benzyl-3-pyrrolidinyl-methyl (4S)-2,6-dimethyl-4-(*m*-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylate hydrochloride, a new calcium antagonist, for 9 wk on blood pressure and urinary excretion of electrolytes were studied in stroke-prone spontaneously hypertensive (SHRSP) rats.
- 2. YM-09730-5 (1 and 3 mg/kg per day, p.o.) prevented development of hypertension and produced a significant reduction in blood pressure from the first week of the experiment. Nicardipine (15 mg/kg per day, p.o.) produced almost the same degree of antihypertensive effect as YM-09730-5 at a dose of 3 mg/kg.
- 3. YM-09730-5 produced significant diuresis and increased urinary excretion of electrolytes throughout the experiment.
- 4. Chronic administration of YM-09730-5 (3 mg/kg) reduced the severity of glomerular lesions in the kidney and vasculitis in the mesenteric artery.
- 5. These results demonstrate that YM-09730-5 is a potential antihypertensive drug with a potency about 5 times higher than that of nicardipine.

<sup>\*</sup> 本報告は Gen. Pharmac. Vol. 22, No. 2, pp. 263-266 (1991) に発表.

<sup>\*\*</sup> 山之内製薬