

Title	Regional differences in extracellular choline dependency of acetylcholine synthesis in the rat brain
Sub Title	
Author	鈴木, 岳之(Suzuki, Takeshi) 鹿島, 裕子(Kashima, Yuko) 藤本, 和子(Fujimoto, Kazuko) 大畑, 尚代(Ohata, Hisayo) 川島, 紘一郎(Kawashima, Koichiro)
Publisher	共立薬科大学
Publication year	1991
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.36 (1991.) ,p.49- 49
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000036-0049

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Regional differences in extracellular choline dependency of acetylcholine synthesis in the rat brain*

Takeshi SUZUKI, Yuko KASHIMA, Kazuko FUJIMOTO, Hisayo OOHATA
and Koichiro KAWASHIMA

鈴木岳之, 鹿島裕子, 藤本和子, 大畑尚代, 川島紘一郎,

Acetylcholine (ACh) synthesis under conditions of restricted extracellular choline uptake was investigated in order to clarify the procurement of choline for ACh synthesis using slices of several regions of the rat brain. Extracellular choline-independent ACh synthesis was observed in the hippocampus, frontal cortex and caudate putamen, which contain cholinergic nerve terminals, whereas little or no synthesis was observed in the medial septum or basal nucleus of Meynert, which contain cholinergic cell bodies. These results indicate that cholinergic nerve terminals, but not the cell bodies, may be able to synthesize choline for ACh biosynthesis. Extracellular choline-dependent ACh synthesis was observed in all regions examined. In the presence of 10 μ M choline, the highest content of newly synthesized ACh and the proportionate increase in ACh compared with the unreleasable fraction (basal level) were observed in the caudate putamen. In the frontal cortex, although the level of synthesized ACh was low, the proportionate increase in ACh was high. In contrast, in the medial septum and basal nucleus of Meynert, high levels of ACh with a low proportionate increase compared with basal levels were observed. In the presence of hemicholinium-3, extracellular choline was also taken up for ACh synthesis in all regions examined, the level being especially high in the frontal cortex and medial septum. The present results indicate that the manner of choline procurement for ACh synthesis is heterogeneous among the various regions of the rat brain.

* 本報告は *Neuroscience Research*, 11 71—76 (1991) に発表.