

Title	Core structure of glycyrrhizan GA, the main polysaccharide from the stolon of glycyrrhiza glabra var. glandulifera ; anti-complementary and alkaline phosphatase-inducing activities of the polysaccharide and its degradation products
Sub Title	
Author	高田, 勝利(Takada, Katsutoshi) 友田, 正司(Tomododa, Masashi) 清水, 訓子(Shimizu, Noriko)
Publisher	共立薬科大学
Publication year	1992
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.37 (1992.) ,p.56- 56
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000037-0056

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Core Structure of Glycyrrhizan GA, the Main Polysaccharide from
the Stolon of *Glycyrrhiza glabra* var. *glandulifera* ;
Anti-Complementary and Alkaline Phosphatase-
Inducing Activities of the Polysaccharide
and Its Degradation Products***

Katsutoshi TAKADA, Masashi TOMODA and Noriko SHIMIZU

高田勝利, 友田正司, 清水訓子

The controlled Smith degradation and limited hydrolysis of glycyrrhizan GA, a representative polysaccharide with remarkable phagocytosis-enhancing activity isolated from the stolon of *Glycyrrhiza glabra* var. *glandulifera* REG. et HERD. were carried out. Methylation analyses of the primary and the secondary Smith degradation products and of the limited hydrolysis product indicated that the core structural features of glycyrrhizan GA include a backbone chain composed of β -1,3-linked D-galactose residues. Three-fifths of the galactose units in the backbone carry side chains composed of β -1,3- and β -1,6-linked D-galactosyl residues at position 6. Anti-complementary and alkaline phosphatase-inducing activities of the polysaccharide, periodate oxidation-reduction and the controlled Smith degradation products were investigated, and the controlled Smith degradation product showed significant activity.

* 本報告は *Chem. Pharm. Bull.*, **40**, 2487—2490 (1992) に発表.