

Title	Practical optimization method for dual-flow countercurrent extraction : purification of glycyrrhizin
Sub Title	
Author	西沢, 秀幸(Nishizawa, Hideyuki) 沖村, 早希子(Okimura, Sakiko) 阿部, 芳広(Abe, Yoshihiro)
Publisher	共立薬科大学
Publication year	1992
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.37 (1992.) ,p.63- 63
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000037-0063

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Practical Optimization Method for Dual-flow Countercurrent Extraction : Purification of Glycyrrhizin*

Hideyuki NISHIZAWA, Sakiko OKIMURA and Yoshihiro ABE

西沢秀幸, 沖村早希子, 阿部芳広

An optimization method for the dual-flow countercurrent extraction of a mixture, the components of which show a concentration dependency in its distribution ratio, has been developed. As an experimental demonstration, commercial glycyrrhizin (about 90% purity) was purified to 99.8% with high recovery using a two-phase system (1-butanol and 0.1M NaCl) in the presence of 0.02M tetra-n-butylammonium hydroxide, and under the optimum conditions of the present method. To acquire 90.0% recovery of glycyrrhizin in an organic extract, the flow rate of the organic solvent was calculated to be 0.943 ml/min (aqueous solvent. 1.06 ml/min). Extraction performed at the derived flow rates obtained 87.2% recovery of glycyrrhizin in the organic extract, showing the propriety of the optimization method.

* 本報告は *Anal. Sci.* 8 (3), 367—374 (1992) に発表.