Title	Large deletion of androsterone UDP-glucuronosyltransferase gene in the inherited deficient strain of wistar rats
Sub Title	
Author	本間, 浩(Honma, Hiroshi) 川合, 博幸(Kawai, Hiroyuki) 久保田, 三保(Kubota, Miho) 松井, 道夫(Matsui, Michio)
Publisher	共立薬科大学
Publication year	1992
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.37 (1992.),p.71- 71
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000037-0071

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Large deletion of androsterone UDP-glucuronosyltransferase
gene in the inherited deficient strain of Wistar rats***

Hiroshi HOMMA, Hiroyuki KAWAI, Miho KUBOTA
and Michio MATSUI

本間 浩, 川合博幸, 久保田三保, 松井道夫

LA Wistar rats have a deficiency of androsterone UDP-glucuronosyltransferase (UDPGT) and are present in Wistar rat colonies around the world. In order to clarify the molecular mechanism of the deficiency, androsterone UDPGT cDNA clone, pGT2 was isolated from rat liver cDNA library and was digested with restriction enzymes to afford 3 probes for Northern and Southern blot analyses in HA (normal), heterozygous LA and LA Wistar rats. In Northern blot analysis, androsterone UDPGT mRNA was totally absent in LA wistar rat liver. Southern blot analysis suggested a large deletion of androsterone UDPGT gene in the rats. Genomic DNA amplifications with synthetic primers which have nucleotide sequences corresponding to the 5'-region of androsterone UDPGT cDNA, suggested that androsterone UDPGT gene has exon I with a length of some 700 bp and that this exon is deleted in LA Wistar rats. Based on these lines of evidence, it is concluded that the large portion of androsterone UDPGT gene is deleted in LA Wistar rats, which results in the absence of androsterone UDPGT mRNA and consequently the corresponding enzyme protein.

* 本報告は *Biochim. Biophys. Acta.*, 1138, 34—40 (1992) に発表.