Title	Immunochemical characterization of developmental changes in rat hepatic hydroxysteroid sulfotransferase
Sub Title	
Author	本間, 浩(Honma, Hiroshi) 中込, 泉(Nakagome, Izumi) 鎌倉, 稔(Kamakura, Minoru) 松井, 道夫(Matsui, Michio)
Publisher	共立薬科大学
Publication year	1992
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.37 (1992.),p.73- 73
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000037-0073

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Immunochemical Characterization of Developmental Changes in Rat Hepatic Hydroxysteroid Sulfotransferase*

Hiroshi HOMMA, Izumi NAKAGOME, Minoru KAMAKURA and Michio MATSUI

本間 浩, 中込 泉, 鎌倉 稔, 松井道夫

A major isoenzyme of hepatic androsterone-sulfating sulfotransferase (AD-ST) was purified from adult female rats. The activity was purified 122-fold over that found in the cytosol and showed a single protein band with a subunit molecular mass of 30 kDa after SDS-polyacrylamide gel electrophoresis. The purified enzyme exhibited four isoelectric variants of subunits on denaturing isoelectrofocusing gels (pIs = 5.8, 6.1, 6.7 and 7.2).

Rabbit antiserum raised against the enzyme specifically detected AD-ST polypeptide in rat liver cytosol. Immunoblot analysis of liver cytosol from female and male rats at various ages showed good correlation between the levels of AD-ST activity and AD-ST polypeptide. Significant levels of AD-ST activity and polypeptide were detected in senescent male rats, though normal adult male rats have very low levels of AD-ST activity and protein. The relative content of the isoelectric variants of AD-ST were different in liver cytosol of weanling and adult females, indicating that age- and gender-related alterations of hepatic AD-ST activity are primarily determined by the levels of AD-ST polypeptide and the relative amounts of the four isoelectric variants of the enzyme.

* 本報告は *Biochim. Biophys. Acta.*, 1121, 69—74 (1992) に発表.