

Title	Substance P-evoked release of acetylcholine from isolated spinal cord of the newborn rat
Sub Title	
Author	小林, 伸行(Kobayashi, Nobuyuki) 佐久間, 雅文(Sakuma, Masafumi) 吉岡, 耕一(Yoshioka, Koichi) 大西, 由子(Onishi, Yuko) 柳澤, 光彦(Yanagisawa, Mitsuhiko) 川島, 紘一郎(Kawashima, Koichiro) 大塚, 正徳(Otsuka, Masanori)
Publisher	共立薬科大学
Publication year	1992
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.37 (1992.) ,p.85- 85
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000037-0085

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

SUBSTANCE P-EVOKED RELEASE OF ACETYLCHOLINE FROM ISOLATED SPINAL CORD OF THE NEWBORN RAT*

N. KOBAYASHI**, M. SAKUMA**, K. YOSHIOKA**, Y. ONISHI**, M. YANAGISAWA**,
K. KAWASHIMA and M. OTSUKA**

小林伸行**, 佐久間雅文**, 吉岡耕一**, 大西由子**, 柳澤光彦**,
川島紘一郎, 大塚正徳**

Isolated spinal cords of newborn rats were perfused with artificial cerebrospinal fluid and the release of endogenous acetylcholine was measured using high-performance liquid chromatography with an electrochemical detection system. Application of high- K^+ (90 mM) medium evoked about an eight-fold increase in the acetylcholine release, and the K^+ -evoked release was Ca^{2+} dependent. Veratridine (20 μ M) also evoked about a four-fold increase in the acetylcholine release, and this increase was suppressed by 0.2 μ M tetrodotoxin. Application of substance P at 0.3–3 μ M evoked a concentration-dependent release of acetylcholine. The substance P-evoked acetylcholine release was Ca^{2+} dependent and abolished by tetrodotoxin. Neurokinin A, neurokinin B, acetyl-Arg⁶-septide and senktide (3 μ M each) also evoked a release of acetylcholine. Electrophysiological experiments using isolated spinal cords of newborn rats showed that bath application of substance P induced a depolarization of motoneurons, which was enhanced by edrophonium. This enhancement of substance P-induced depolarization by edrophonium disappeared in a low- Ca^{2+} medium or in the presence of atropine and dihydro- β -erythroidine. In the presence of edrophonium and dihydro- β -erythroidine, substance P induced an inhibition of monosynaptic reflex, and this inhibition was abolished by atropine.

These results suggest that substance P and other tachykinins induce a release of acetylcholine from the newborn rat spinal cord by exciting cholinergic neurons.

* 本報告は *Neuroscience* vol. 45, No. 2, pp. 331–337. 1991 に発表.

** 東京医科歯科大学 医学部薬理学教室