

Title	Acetylcholine synthesis is modulated by acetylcholine content of cytosolic fraction but not by that of releasable fraction
Sub Title	
Author	鈴木, 岳之(Suzuki, Takeshi) 鹿島, 裕子(Kashima, Yuko) 藤本, 和子(Fujimoto, Kazuko) 川島, 紘一郎(Kawashima, Koichiro)
Publisher	共立薬科大学
Publication year	1992
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.37 (1992. ) ,p.86- 86
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	<a href="https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000037-0086">https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000037-0086</a>

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Acetylcholine synthesis is modulated by acetylcholine  
content of cytosolic fraction but not by that of  
releasable fraction\***

Takeshi SUZUKI, Yuko KASHIMA, Kazuko FUJIMOTO  
and Koichiro KAWASHIMA

鈴木岳之, 鹿島裕子, 藤本和子, 川島紘一郎

Synthesis and release of acetylcholine (ACh) in the rat hippocampal slices were examined to clarify the mechanism of modulation of ACh synthesis. Treatment with 2-(4-phenylpiperidino)cyclohexanol (AH 5183, 50  $\mu$ M), an inhibitor of ACh transport from cytosol to synaptic vesicles, inhibited the increase in ACh content of the membrane-bound fraction which is readily releasable, but did not affect the cytosolic ACh content. Under these conditions. The total ACh content reached a plateau value. These results indicate that ACh synthesis is modulated by cytosolic ACh content but not by the vesicular fraction.

---

\* 本報告は *Neuroscience Letters*, 144 (1992) 127—129 に発表.