Title	Regional assignment of rat androsterone UDP-glucuronosyltransferase gene (UGT2B2) to chromosome 14p21.2-p22
Sub Title	
Author	佐藤, 均(Sato, Hitoshi) 永井, 総子(Nagai, Fusako) 本間, 浩(Honma, Hiroshi) 森, 茂郎(Mori, Shigeo) 松井, 道夫(Matsui, Michio)
Publisher	共立薬科大学
Publication year	1993
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.38 (1993.),p.41- 41
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000038-0041

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Regional Assignment of Rat Androsterone UDP-glucuronosyl-transferase Gene (UGT2B2) to Chromosome 14 p21.2-p22*

Hitoshi SATOH**, Fusako NAGAI, Hiroshi HOMMA, Sigeo MORI**, and Michio MATUI

佐藤 均**, 永井総子, 本間 浩, 森 茂郎**, 松井道夫

UDP-glucuronosyltransferase (UGT) consists of a multigene family and there appear to be more than eight UGT isoenzymes in the rat. Rat liver androsterone UGT (UGT2B-2) catalyzes the glucuronidation of androsterone, etiocholanolone, lithocholic acid and naphthylamines. The whole construct of the recombinant plasmid pGT2 containing UGT2B2 cDNA was used as a probe. The DNA probe was labeled by nick-translation with biotin-16-dUTP. Prometaphase chromosomes were prepared from primary lung fibroblast cultures of male Wistar rats. Chromosome slides were prepared, hybridized with the DNA probe and incubated with fluorescein avidin DCS. The intensity of the fluorescence was amplified by addition of a layer of biotinylated goat anti-avidin D and the slides were stained with propidium iodide. The signals were stained as yellow-green spots on red chromosomes and the location of the signals were recognized on Q-banded chromosomes. In the present study, we regionally assigned UGT2B2 gene to chromosome 14p21.2-p22. UGT2B genes have been mapped to human chromosome 4 and mouse chromosome 5, respectively. It is known that the albumin and α -fetoprotein genes are closely linked on human chromosome 4, mouse chromosome 5 and rat chromosome 14. These results indicate that there remains a certain degree of linkage homology in these genes among human, mouse and rat.

* 本報告は *Cytogenet. Cell Genet.*, **62**, 49—51 (1993) に発表.

** 東大医科研・病理学研究部