

Title	Arabinogalactan core structure and immunological activities of ukonan C, an acidic polysaccharide from the rhizome of curcuma longa
Sub Title	
Author	権田, 良子(Gonda, Ryoko) 友田, 正司(Tomododa, Masashi) 大原, 直子(Ohara, Naoko) 高田, 勝利(Takada, Katsutoshi)
Publisher	共立薬科大学
Publication year	1993
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.38 (1993.) ,p.45- 45
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000038-0045

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Arabinogalactan Core Structure and Immunological Activities
of Ukonan C, an Acidic Polysaccharide from
the Rhizome of *Curcuma longa****

Ryōko GONDA, Masashi TOMODA, Naoko ŌHARA and Katsutoshi TAKADA

権田良子, 友田正司, 大原直子, 高田勝利

Controlled Smith degradation of ukonan C, a phagocytosis-activating polysaccharide isolated from the rhizome of *Curcuma longa* L., was performed. The reticuloendothelial system-potentiating, anti-complementary and alkaline phosphatase-inducing activities of ukonan C and its degradation products were investigated. Methylation analyses of the primary and secondary Smith degradation products and of a de-arabinosylated product indicated that structural features of the arabinogalactan core of ukonan C include a backbone chain composed of β -1,3-linked D-galactose and β -1,4-linked D-xylose. All of the galactose units in the backbone carry side chains composed of β -1,6-linked D-galactosyl residues with or without terminal α -L-arabinose units at position 3.

Ukonan C showed remarkable effects on both reticuloendothelial system-potentiating and alkaline phosphatase-inducing activities. Periodate oxidation caused a decrease in or disappearance of the immunological activities, but the controlled Smith degradation product having the arabinogalactan core structure of polysaccharide showed a pronounced effect on anti-complementary activity.

* 本報告は *Biol. Pharm. Bull.*, **16** (3), 235—238 (1993) に発表.