

Title	Characterization of two novel polysaccharides having immunological activities from the root of panax ginseng
Sub Title	
Author	友田, 正司(Tomoda, Masashi) 平林, 啓子(Hirabayashi, Keiko) 清水, 訓子(Shimizu, Noriko) 権田, 良子(Gonda, Ryoko) 大原, 直子(Ohara, Naoko) 高田, 勝利(Takada, Katsutoshi)
Publisher	共立薬科大学
Publication year	1993
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.38 (1993.) ,p.47- 47
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000038-0047

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Characterization of Two Novel Polysaccharides Having
Immunological Activities from the Root
of *Panax ginseng****

Masashi TOMODA, Keiko HIRABAYASHI, Noriko SHIMIZU, Ryōko GONDA,
Naoko ŌHARA and Katsutoshi TAKADA

友田正司, 平林啓子, 清水訓子, 権田良子, 大原直子, 高田勝利

Two acidic polysaccharides, called ginsenan S-IA and ginsenan S-IIA, were isolated from the root of *Panax ginseng* C. A. MEYER. They were homogeneous on electrophoresis and gel chromatography, and their molecular masses were estimated to be 5.6×10^4 , and 1.0×10^5 , respectively. Ginsenan S-IA is composed of L-arabinose : D-galactose : D-galacturonic acid in the molar ratio of 8 : 8 : 1, and ginsenan S-IIA is composed of L-arabinose : D-galactose : D-glucose : D-galacturonic acid in the molar ratio of 15 : 10 : 2 : 5, in addition to small amounts of O-acetyl groups. About a half (ginsenan S-IA) and about a quarter (ginsenan S-IIA) of the hexuronic acid residues exist as methyl esters. Reduction of carboxyl groups, methylation analysis, nuclear magnetic resonance and periodate oxidation studies indicated that their structural features include mainly α -1,5-linked L-arabino- β -3,6-branched D-galactan type structural units. Both polysaccharides showed remarkable reticuloendothelial system-potentiating activity in a carbon clearance test and pronounced anti-complementary activity. These substances are the first examples having a relatively high content of both α -3,5-branched L-arabinose and β -1,4-linked D-galactose units among the acidic arabinogalactans with activities on phagocytosis and anti-complement.

* 本報告は *Biol. Pharm. Bull.*, **16** (11), 1087—1090 (1993) に発表.