Title	Sterol metabolism in insects : dealkylation of phytosterol to cholesterol
Sub Title	
Author	池川, 信夫(Ikekawa, Nobuo) 森崎, 益雄(Morisaki, Masuo) 藤本, 善徳(Fujimoto, Yoshinori)
Publisher	共立薬科大学
Publication year	1993
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.38 (1993.) ,p.64- 64
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000038-0064

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Sterol Metabolism in Insects : Dealkylation of Phytosterol to Cholesterol

Nobuo IKEKAWA**, Masuo MORISAKI and Yoshinori FUJIMOTO***

池川信夫、森崎益雄、藤本善徳

Insects obtain indispensable sterol by the modification of C-24-alkylated dietary sterols. Since this process is extremely important for insects, they have employed a diversified enzyme system with an excellent ability to dealkylate a variety of phytosterols, regardless of whether the C-24-alkyl group is methyl or ethyl, whether the stereochemistry at C-24 is *R* or *S*, and whether the C-22 double bond is present or absent. The dealkylation of phytosterol is carried out by three successive reactions: dehydrogenation, epoxidation, and epoxide fragmentation. The resulting desmosterol is eventually hydrogenated to cholesterol. The crucial step is fragmentation of epoxide, and this reaction appears to occur in a highly stereoselective manner, as evidenced from the *pro-S*- and *pro-R*-methyl groups of fucosterol (24*R*, 28*R*)-epoxide being transformation to the (*Z*)- and (*E*)-methyl groups, respectively, of desmosterol. Experimental evidences of all these results obtained by the use of the silkworm *Bombyx mori* have been presented. Also discussed are sterol structure requirement of *B. mori*, and inhibitors of sterol metabolism in insects.

* 本報告は *Acc. Chem. Res.*, **26**, 139—146, (1993) に発表。

** いわき明星大学・理工学部

*** 東京工業大学・理学部