慶應義塾大学学術情報リポジトリ

Keio Associated Repository of Academic resouces

New Associated Repository of Academic resources	
Title	Aerial oxidation of some 2-pyridyl ketone hydrazones catalyzed by Cu ²⁺ . physical properties of reaction products
Sub Title	
Author	森, 久和(Mori, Hisakazu) 坂本, 香織(Sakamoto, Kaori) 増戸, 幸枝(Mashito, Sachie) 松岡, 陽子(Matsuoka, Yoko) 松林, 美佳(Matsubayashi, Mika) 酒井, 和恵(Sakai, Kazue)
Publisher	共立薬科大学
Publication year	1993
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.38 (1993.),p.69-69
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000038-0069

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Aerial Oxidation of Some 2-Pyridyl Ketone Hydrazones Catalyzed by Cu²⁺. Physical Properties of Reaction Products*

Hisakazu Mori, Kaori Sakamoto, Sachie Mashito, Yohko Matsuoka, Mika Matsubayashi and Kazue Sakai

森 久和,坂本香織,増戸幸枝,松岡陽子,松林美佳,酒井和恵

Some 2-pyridyl ketone hydrazones were subjected to aerial oxidation catalyzed by Cu²⁺. The oxidation product of di-2-pyridyl ketone hydrazone had the structure of a derivative of [1,2,3]-triazolo[1,5-a] pyridine, and this structure is different from that previously proposed by other investigators. The fluorescence spectra of the oxidation products were measured in solutions of a wide pH range. In contrast to the oxidation products of other 2-pyridyl ketone hydrazones, that of di-2-pyridyl ketone hydrazone showed very strong fluorescence in acidic media. The characteristic nature of this compound was also apparent in the ultraviolet spectrum. The generation of hydroxyl radical was demonstrated in the aerial oxidation of di-2-pyridyl ketone hydrazone catalyzed by Cu²⁺, suggesting the formation of hydrogen peroxide as another oxidation product.

^{*} 本報告は Chem. Pharm. Bull. 41 (11), 1944—1947 (1993) に発表。