

Title	[ <sup>3</sup> H] 2-(4-phenylpiperidino) cyclohexanol (AH 5183) binding to synaptosomes and subcellular fractions obtained from rat brain
Sub Title	
Author	鈴木, 岳之(Suzuki, Takeshi) 鹿島, 裕子(Kashima, Yuko) 藤本, 和子(Fujimoto, Kazuko) 川島, 紘一郎(Kawashima, Koichiro)
Publisher	共立薬科大学
Publication year	1993
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.38 (1993. ),p.77- 77
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	<a href="https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000038-0077">https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000038-0077</a>

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**[<sup>3</sup>H]2-(4-Phenylpiperidino)cyclohexanol (AH 5183) binding  
to Synaptosomes and Subcellular Fractions Obtained  
from Rat Brain\***

Takeshi SUZUKI, Yuko KASHIMA, Kazuko FUJIMOTO and Koichiro KAWASHIMA

鈴木岳之, 鹿島裕子, 藤本和子, 川島紘一郎

We investigated the binding of [<sup>3</sup>H]AH 5183 (2-(4-phenylpiperidino)cyclohexanol) to rat brain synaptosomes and subcellular fractions. A high content of specific binding was observed in crude synaptosomes obtained from the striatum, cerebral cortex and hippocampus. The highest density of subcellular binding sites was observed in the synaptic vesicle-rich fraction. The affinities of AH 5183 binding to crude synaptosomes and the synaptic vesicle-rich fraction were almost equivalent, but the density of binding sites was higher in the synaptic vesicle fraction. The present findings indicate that [<sup>3</sup>H]AH 5183 binding to both synaptosomes and the synaptic vesicle-rich fraction is useful as a cholinergic marker, and that for quantitative studies. binding to the latter fraction is more preferable.

---

\* 本報告は *Neuroscience Letters*, 157 : 72-74 (1993) に発表。