Title	Effects of 2-mercaptopropionylglycine on the development of X-ray-induced cataract in rats
Sub Title	
Author	小林, 静子(Kobayashi, Shizuko) 粕谷, 美南子(Kasuya, Minako) 石井, 康雄(Ishii, Yasuo) 竹鼻, 眞(Takehana, Makoto) 酒井, 一夫(Sakai, Kazuo) 鈴木, 紀夫(Suzuki, Norio) 糸井, 素一(Itoi, Motokazu)
Publisher	共立薬科大学
Publication year	1993
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.38 (1993.),p.78- 78
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000038-0078

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Effect of 2-Mercaptopropionylglycine on the Development of X-ray-induced Cataract in Rats

Shizuko KOBAYASHI, Minako KASUYA**, Yasuo ISHII**, Makoto TAKEHANA,
Kazuo SAKAI***, Norio SUZUKI*** and Motokazu ITOI**

小林静子, 粕谷美南子**, 石井康雄**, 竹鼻 眞, 酒井一夫***,
鈴木紀夫***, 糸井素一**

The effect of 2-mercaptpropionylglycine on the development of cataract induced by a single dose of X-ray (10 Gy) was investigated in rats. Intraperitoneal injection of 20 mg/kg, three times weekly starting 1 day after irradiation delayed the development of X-ray-induced cataracts significantly. The amounts of non-protein SH groups, malondialdehyde and the Na^+/K^+ ratio, in the lenses of rats post-treated with the drug were significantly maintained at normal levels even at 27 weeks after irradiation.

On the other hand, a single administration of 250 mg/kg of the compound, 30 min prior to irradiation had no effect on cataract progression induced by X-ray.

* 本報告は *Curr. Eye Res.*, 11, 1099—1103 (1992) に発表。

** 白内障研究所

*** 東京大学医学部