

Title	The actual state of the physically handicapped students in physical education at Keio University
Sub Title	
Author	依田, 隆也(Yoda, Ryuya)
Publisher	慶應義塾大学体育研究所
Publication year	1974
Jtitle	体育研究所紀要 (Bulletin of the institute of physical education, Keio university). Vol.14, No.1 (1974. 12) ,p.69- 69
JaLC DOI	
Abstract	
Notes	Abstract
Genre	
URL	<a href="https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00135710-00140001-0069">https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00135710-00140001-0069</a>

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

## The Actual State of the Physically Handicapped Students in Physical Education at Keio University

By *Ryuya Yoda*\*

This is a report on the study of the actual state of the physically handicapped students who received physical education at Keio University during the past five years. The following is the result obtained.

1. Those students who were judged to require a limit to the load of the physical training were nearly four per cent of the total number of students both in males and females.
2. As to disorders in the kinetic organs, the disorders of the spine such as hernia of intervertebral disc were most frequent in male students, but only a few female students had such disorders.
3. In the students with past history of nephritis, the rate of urinary protein positive reaction was remarkably high. It was assumed that not a small number of them were in the stage of chronic nephritis or recurrent nephritis.
4. As to disorders of the circulatory organs, a relatively large number of male students had hypertension, while in many female students, anemia and hypotension were found.
5. As to disorders of the respiratory organs, the frequently encountered condition was the past history of spontaneous pneumothorax in male students, however, such a case history was found rarely in female students.

---

\* Professor of the Institute of Physical Education, Keio University.