

Title	Serum lactate dehydrogenase (LDH) and prolonged exercise : LDH-1 and possible heart muscle injuries after prolonged exercise
Sub Title	
Author	橋本, 治雄(Hashimoto, Haruo) 中村, 毅志夫(Nakamura, Kishio) 近藤, 明彦(Kondo, Akihiko) 松田, 雅之(Matsuda, Masayuki)
Publisher	慶應義塾大学体育研究所
Publication year	1982
Jtitle	体育研究所紀要 (Bulletin of the institute of physical education, Keio university). Vol.22, No.1 (1982. 12) ,p.81- 82
JaLC DOI	
Abstract	
Notes	Abstract
Genre	
URL	<a href="https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00135710-00220001-0081">https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00135710-00220001-0081</a>

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

# Serum Lactate Dehydrogenase (LDH) and Prolonged Exercise

— LDH-1 and Possible Heart Muscle Injuries  
after Prolonged Exercise —

By *Haruo Hashimoto\**  
*Kishio Nakamura\*\**  
*Akihiko Kondoh\*\*\**  
*Masayuki Matsuda\*\*\*\**

Prolonged swimming in the fresh water pool were chosen for the investigation of the influence of prolonged exercise on the changes in the blood Lactate Dehydrogenase (LDH) levels.

Fifteen students of the varsity swimming team at KEIO University performed prolonged swimming on July 5, 1982 as one of the preparatory training for the channel swimming between Ito and Shimoda (Shizuoka-ken) which was executed on July 27, 1982. Four subjects swam for 6 hours 30 minutes (6:30). Five subjects swam for 7 hours 30 minutes (7:30). Six subjects swam for 9 hours 30 minutes (9:30).

Five students of the varsity track and field team performed 1500 m and/or 5000 m races on July 5, 1982 which were chosen for the comparison of shorter time exercise.

Serum LDH were determined before and after the prolonged exercise (swimming) and track and field events. Serum LDH rose moderately after the 6:30 and 7:30 hours prolonged swimming, while that of the 9:30 hours swimming rose remarkably.

Isoenzymatic analysis revealed that most of the total LDH increment were due to increase of the LDH-5 and LDH-4 which originated from skeletal muscle. LDH-1 which originated from cardiac muscle increased significantly only in the subjects of 9:30 hours prolonged swimming. These results suggests that there may be a cardiac

---

\* Associate Professor of the Institute of Physical Education, Keio University. (M.D.)  
\*\* Assistant Professor of the Institute of Physical Education, Keio University. (M.D.)  
\*\*\* Assistant of the Institute of Physical Education, Keio University.  
\*\*\*\* Assistant of the Institute of Physical Education, Keio University.

muscle injury in the subjects of 9:30 hours prolonged swimming.

LDH isoenzymatic analysis is very useful methods to detect the presence of cardiac muscle injury in the training subjects.