

Title	Discussion of the center of gravity sway of standing posture : its relation to the continuation of physical exercise
Sub Title	
Author	佐々木, 三男(Sasaki, Mitsuo)
Publisher	慶應義塾大学体育研究所
Publication year	1984
Jtitle	体育研究所紀要 (Bulletin of the institute of physical education, Keio university). Vol.24, No.1 (1984. 12) ,p.93- 94
JaLC DOI	
Abstract	
Notes	Abstract
Genre	
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00135710-00240001-0093

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Discussion of the Center of Gravity Sway of Standing Posture—Its Relation to the Continuation of Physical Exercise

By *Mitsuo Sasaki**

Is it possible that body balance, which is one of the common features to superior sports competitors, can be improved by the continuation of physical exercise?

The purpose of this study is to provide basic materials for body balance required in practicing exercise.

The method used in the experiment was to measure the length of sway (mm) and the area of sway (cm²) in standing posture while subjects opened or closed their eyes for 20 seconds respectively, as well as while they stood only on their left legs or right legs for 10 seconds.

The equipment used in the measurement were the Gravicorder (Model G4301S made by Anima) and X.Y. Recorder (Model G1800 made by Anima).

The subjects participated in the experiment were (1) ninety-two healthy untrained males, (2) seventeen male athletes, twelve male basketball players and eight male gymnasts who were training themselves to participate in extramurals, and (3) eleven basketball players representing Japan.

Results and Discussion

(1) In standing posture, the length of sway can somewhat be reduced by the continuation of physical exercise.

(2) As to the length of sway and the area of sway, it seems that biomechanical meaning of the former differs from that of the latter. That is, in the study of this time, the length of sway was reduced by the continuation of exercise, however, no change was noted in the area of sway, while the eyes were opened.

(3) In the case of standing with one leg only, nothing superior was noted in either leg of the subjects who have continued the physical exercise compared with

* Assistant Professor of the Institute of Physical Education, Keio University.

who not continued.

(4) The center of gravity on foot was found to be located close to the toe by the continuation of exercise.