

Title	On the production of antibiotics (II) studies on n-butylthiomethyl-penicillin
Sub Title	
Author	須網, 哲夫(Suami, Tetsuo)
Publisher	慶應義塾大学藤原記念工学部
Publication year	1951
Jtitle	Proceedings of the Fujihara Memorial Faculty of Engineering Keio University Vol.4, No.14 (1951.) ,p.79(19)- 85(25)
JaLC DOI	
Abstract	The recent eminent publication by Behrens et al on the biosynthesis of new penicillins has made the present author to investigate the n-butylthiomethyl-penicillin from the industrial point of view. The author studies the biosynthetic effects of n-butylmercaptoacetic acid on shaking culture using <i>Penicillium chrysogenum</i> Q 176 pigmentless saltant. The n-butylthiomethyl-penicillin sodium salt is extracted by solvent extraction and purified by chromatography. Using the pure sodium salt of n-butylthiomethyl-penicillin prepared, the stabilities and the effective distribution coefficients between water and organic solvents are estimated.
Notes	
Genre	Departmental Bulletin Paper
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=KO50001004-00040014-0019

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

On the Production of Antibiotics (II)

Studies on n-Butylthiomethyl-penicillin*

(Received May 18, 1950)

Tetsuo SUAMI**

Abstract

The recent eminent publication by Behrens et al on the biosynthesis of new penicillins has made the present author to investigate the n-butylthiomethyl-penicillin from the industrial point of view.

The author studies the biosynthetic effects of n-butylmercaptoacetic acid on shaking culture using *Penicillium chrysogenum* Q 176 pigmentless saltant. The n-butylthiomethyl-penicillin sodium salt is extracted by solvent extraction and purified by chromatography. Using the pure sodium salt of n-butylthiomethyl-penicillin prepared, the stabilities and the effective distribution coefficients between water and organic solvents are estimated.

I. Introduction

The recent eminent publication by Behrens et al¹⁾ on the biosynthesis of new penicillins has made the present author to investigate the n-butylthiomethyl-penicillin from the industrial point of view.

n-Butylmercaptoacetic acid is prepared by Larsson's method²⁾ from thioglycolic acid and n-butylbromide and at this juncture the author find that this condensation proceeds more smoothly and rapidly (5 hrs.) in a methanol-water (3:1) solution, yield 62.8%.

The author studies the biosynthetic effects of n-butylmercaptoacetic acid on shaking culture using *Penicillium chrysogenum* Q 176 pigmentless saltant. The n-butylthiomethyl-penicillin sodium salt is extracted by solvent extraction and purified by chromatography. Using the pure sodium salt of n-butylthiomethyl-penicillin prepared, the stabilities and the effective distribution coefficients between water and three various organic solvents (butylacetate, ethylether and chloroform) are estimated.

I wish to express my sincere and profound gratitude to Prof. Dr. Sumio Umezawa for his continuous and kind guidance. I also extend my heartfelt gratitude to Mr. Masao Naito who kindly helped me in carry out the experiments.

* This report was printed in *Journal of Antibiotics* 4 (1) 9 (1950) (in Japanese).

** Assistant Professor at Keio University.

1) *J. Biol. Chem.*, 175, 771 (1948)


2) *Larsson, Ber.*, 63, 1340 (1930)

II. Experiments

1. n-Butylmercaptoacetic acid

(a) Thioglycolic acid

Three reagents, namely, potassium hydrosulfide³⁾, sodium hydrosulfide and sodium polysulfide are reacted with monochloroacetic acid⁴⁾ respectively. The author find that the first reagent and the second give lower yield (about 45% yield) and the third comperatively better yield. (62.8% yield).



Hot sodium disulfide which is prepared by the addition of 6 gr. of powdered sulfur to 45 gr. of sodium sulfide ($\text{Na}_2\text{S} \cdot 9\text{H}_2\text{O}$) fused in a procelain crucible, is added into the mixed solution of 40 gr. of monochloroacetic acid, and saturated water solution of 17 gr. of sodium hydroxide. The temperature of the reaction mixture rises and the reaction finishes in few minutes. The reaction mixture is added slowly with 160 gr. of 50% sulfuric acid and filtered. 30 gr. of zinc tablets are added to the filtrate with agitation and after zinc all disappears, the reaction mixture is filtered to remove zinc sulfate. The filtrate is extracted two times with 50 cc ether, and the ether layer dried over sodium sulfate. The solvent is removed under reduced pressure and residual oil fractionally distilled in vacuo. Thioglycolic acid 27 gr. is obtained, b. p. $123^\circ\text{C}/27\text{mm}$ ($107-108^\circ\text{C}/16\text{mm}$), yield 70%.

(b) n-Butylbromide

n-Butylbromide is prepared by the method described in Organic Synthesis Vol. I, 28.

(c) n-Butylmercaptoacetic acid

The auther prepares n-butylmercaptoacetic acid by the method of Larsson²⁾ with 40% yield and tries a modified method, obtaining better yield.

The mixed solution of methanol 300 cc, distilled water 100 cc, sodium hydroxide 20 gr. and thioglycolic acid 25 gr. is added with n-butylbromide 36 gr. under continuous agitation. After 5 hrs. methanol is removed from the reaction mixture by distillation and residual oil is washed with small portions of ether. This is acidified by 20% sulfuric acid and extracted two times with 50 cc ether. The combined ether-extract is dried over sodium sulfate, and ether evaporated under reduced pressure. n-Butylmercaptoacetic acid a colourless 45.2 gr. viscous oil is obtained by distillating the residue in vacuo. b. p. $152^\circ\text{C}/20\text{mm}$ yield 62.8%.

2. Submerged fermentation with n-butylmercaptoacetic acid

(a) With 0.018% n-butylmercaptoacetic acid

Penicillium chrysogenum Q 176 pigmentless saltant is inoculated into each flask

3) Klason and Carson, Ber. 39, 732 (1906)

4) Friedlander, Monatsheft f. Chem. 28, 247

of 500 cc volume containing 200 cc of corn steep medium (corn steep 4.0%, lactose 3.0%, glucose 0.5%, calcium carbonate 1.0%, NaHPO_4 0.15%, urea 0.1%, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.025%, $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ 0.00056%, n-butylmercaptoacetic acid 0.018%, pH 6.5). They are cultivated at $24^\circ \pm 1^\circ\text{C}$ on the rotatory shaking machine (150 r. p. m.). On the other hand, the medium without n-butylmercaptoacetic acid is used as the control. The potency is measured by the cylinder-plate method according to F. D. A. penicillin assay method using F. D. A. 209 p as the test organism. The results are shown in Fig. 1.




Fig. 1. Potency increase in the medium with 0.018% n-butylmercaptoacetic acid




Fig. 2. Influence of the concentration of n-butylmercaptoacetic acid on potency

(b) Optimum concentration of n-butylmercaptoacetic acid

To the same medium as (a), 0.04%, 0.05% and 0.06% of n-butylmercaptoacetic acid is added respectively. *Penicillium chrysogenum* Q 176 pigmentless saltant is used. Fermentation is done in the same conditions as (a). The medium with 0.3% phenylacetic acid instead of n-butylmercaptoacetic acid is used as the control. The results are shown in Fig. 2.

3. Sodium salt of n-butylthiomethyl-penicillin

A 10 L fermentation broth obtained as described in 2 (a) (fermentation 120 hrs., potency 970 u/cc) is cooled with ice, acidified by 10% sulfuric acid and extracted two times with 3.0 L amylacetate at pH 2.0. The amylacetate solution is extract with 2.0 L phosphate-buffer solution (pH 7.0) and then 2.1 L of the first rich-water (potency 4,000 u/cc, total potency 8,400.000 u) is obtained. This first rich-water is extracted with 0.6 L amylacetate two times repeatedly at 0°C , pH

2.0 and total amyacetate solution is titrated with freshly made 5.0% sodium bicarbonate solution to pH 6.8. A second rich water 93 cc (potency 96,000 u/cc, total potency 800,000 u.,) is obtained. The combined yield is 82%.

200 cc of the second rich-water is extracted with 30 cc of amyacetate two times at 0°C, pH 2.0 and combined amyacetate layer is dried over sodium sulfate. The dried amyacetate solution is percolated through the column (diameter 2.1 cm, length 20 cm) of 30 gr. active alumina BL 6 of "Nippon Aluminium Co., Ltd." and then the development is done with fresh butylacetate. The column is cutted from the top in eight fractions of 2.5 cm length and each fraction extracted with 50 cc of phosphate-buffer solution (pH 7.0). The results are shown in Table 1.

Table 1

Fraction	Units in Fractions	%	Differential Assay Value*
1	449,000	33.0	0.64
2	429,000	31.5	0.54
3	295,000	21.7	0.53
4	122,000	9.0	—
5	39,000	2.9	—
6	29,300	2.5	—
7	None	0	—
8	None	0	—
Total	1,363,300		

The mixture of the second and the third fraction is extracted with 50 cc butylacetate two times at 0°C, pH 2.0 and total butylacetate is titrated to pH 6.8 with fresh 5% sodium bicarbonate solution.

The final rich-water (110,000 u/cc) 6.2 cc is dried in the frozen state and n-butylthiomethyl-penicillin sodium salt (290 mg, 2,350 u/mg) is obtained.

This is dissolved in 1.5 cc of acetone-water solution (9:1) and after the further addition of 5.0 cc absolute acetone, the solution is cooled in ice. A crystalline sodium salt of n-butylthiomethyl-penicillin salt (200 mg, 2,980 u/mg) is recovered. The potency is elevated by repeated recrystallization (140 mg, 3,310 u/mg). The sodium salt of crystalline n-butylthiomethyl-penicillin represents 0.54 as the differential assay value.

4. Procain salt of n-butylthiomethyl-penicillin

20 cc of the final rich-water described above is added with 1.8 cc of the aqueous solution of 50% procain hydrochloride under agitation and the mixture is stored in refrigerator over night. Prisms are obtained. (potency 1,800 u/mg, yield 803 mg, 84%).

* Test organisms are *B. subtilis* N. R. R. L. 558 (R) and *St. aureus* 209 P.

After thrice recrystallization with methanol, n-butylthiomethyl-penicillin procain salt (510 mg) of m. p. 110°C, and potency 2,100 u/mg, is obtained.

5. Calcium salt of n-butylthiomethyl-penicillin

210 mg sodium salt of n-butylthiomethyl-penicillin (potency 3,310 u/mg, total potency 695,000 u.) is dissolved in 10 cc sterilized distilled water at 0°C and extracted three times with 10 cc butylacetate at pH 2.5. After total butylacetate layer is washed with small portions of chilled water and 200 mg of sedimentary calcium carbonate are added to the butylacetate layer and the mixture are shaken for ten minutes. The aqueous layer which is separated by the centrifugal separator from the mixed solution, is filtered through the Seitz filter and 4.0 cc of the final solution of n-butylthiomethyl-penicillin calcium salt (potency 130,000 u/cc, total potency 520,000 u, yield 74.8%) is obtained. On drying in the frozen state, 159 mg of calcium salt of n-butylthiomethyl-penicillin (potency 3,230 u/mg, total potency 515,000 u, yield 74%) are obtained. This salt has one molecular water of crystallisation and after drying at 60°C in vacuo, calcium is determined.

Sample : 6,920 mg, drying loss : 0.160 mg, Calcd, for

$C_{14}H_{21}O_4N_2S_2 \cdot \frac{1}{2}Ca \cdot \frac{1}{2}H_2O : H_2O, 2.41 \cdot$ Found $H_2O, 2.31$

Sample : 4.821 mg, 5.423 mg ; $GaSO_4 : 0.904$ mg, 1.009 mg ;

Calcd. for $C_{14}H_{21}O_4N_2S_2 \cdot \frac{1}{2}Ca : Ca$ 5.49

Found. Ca : 5.49 5.48

6. Stabilities of n-butylthiomethyl-penicillin in aqueous solution

20 cc of sterilized distilled water in which 13 mg of n-butylthiomethyl-penicillin sodium salt (3,310 u/mg) is dissolved is adjusted to pH 2.0 at 0°C, with 10% sulfuric acid. Maintaining this solution at 0°C, 1.0 cc is taken up from this solution at every interval diluted with phosphate buffer solution (pH 6.5) and immediately assayed by cylinder-plate method. The mean half-value period of n-butylthiomethyl-penicillin is calculated. Results are shown in Table 2 No. 1.

100 cc of sterilized distilled water in which 8.5 mg sodium salt of n-butylthiomethyl-penicillin (3,310 u/mg) is dissolved is adjusted to pH 2.5, with 10% sulfuric acid at 0°C. The mean half-value period is calculated as shown in Table 2 No. 2.

10 cc of sterilized phosphate-buffer solution (pH 6.5) in which 6.3 mg sodium salt of n-butylthiomethyl-penicillin (3,310 u/mg) are dissolved are maintained at 0°C and assaying this solution at intervals, the mean half-value period shown in Table 2 No. 3 is calculated.

100 cc of sterilized phosphate-buffer solution (pH 6.5) in which 8.5 mg sodium salt of n-butylthiomethyl-penicillin (3,310 u/mg) are dissolved are maintained at 0°C and assaying this solution at intervals, the mean half-value period shown in Table 2 No. 4, is calculated.

Table 2

Hrs	No. 1 pH 2.0		No. 2 pH 2.5		No. 3 pH 6.5		No. 4 pH 6.5	
	u/cc	k × 10	u/cc	k × 10	u/cc	k × 10 ²	u/cc	k × 10 ²
0.0	2,090		270		2,080		281	
1.0	1,800	1.5	—		—		—	
3.0	1,370	1.4	—		—		—	
6.0	—		228	0.28	—		—	
7.0	—		—		2,030	0.44	—	
8.0	660	1.4	—		—		—	
10.0	487	1.4	—		—		—	
12.0	—		193	0.28	1,990	0.43	—	
22.0	—		144	0.29	—		270	0.18
48.0	—		—		—		261	0.15
Mean	K = 1.4 × 10 ⁻¹		K = 0.28 × 10 ⁻¹		K = 0.43 × 10 ⁻²		K = 0.17 × 10 ⁻²	
Half-value period (hrs)	4.9		24.7		157		408	

7. The effective distribution coefficients

5.0 cc of the aqueous solution of various concentration of the n-butylthiomethylpenicillin sodium salt (3,310 u/mg) is added with 5.0 cc of organic solvent. The mixture is taken vigorously at 0°C, adjusted to pH 2.5 with 10% phosphoric acid and then aqueous layer is separated by a centrifugal separator. The aqueous layer is diluted with phosphate-buffer solution (pH 6.5) and the effective distribution coefficients are calculated from the potency of the diluted solution. Results are shown in Table 3.

Table 3

Solvents	Potency u/cc			Distribution coefficient
	Initial	Water layer	Organic sol- vent layer	
Butylacetate	996	16.0	980	61.3/1
"	204	3.8	200	52.6/1
Ether	996	26.0	970	37.3/1
"	238	6.1	232	38.0/1
Chloroform	204	10.7	193	18.4/1
"	238	19.2	219	11.4/1

III. Summary

(1) n-Butylmercaptoacetic acid is prepared by the method of Larsson²⁾ from thioglychic acid and n-butylbromide, and at this juncture the author find that this condensation proceeds more smoothly and rapidly (5 hrs.) in a methanol-water (3 : 1) solution, yield 62.8%.

(2) The optimum concentration of n-butylmercaptoacetic acid in culture medium on submerged fermentation using *Penicillium chrysogenum* Q 176 pigmentless saltant is about 0.05%, and at this condition, a stimulation ratio is 2.46.

(3) The sodium salt of n-butylthiomethyl-penicillin (3,310 u/mg) is obtained by chromatographic purification of crude products which is separated by solvent extraction from the fermentation broth. The sodium salt is converted into the calcium salt and the analysis of calcium salt gives a satisfactory result.

(4) From the final rich-water, the procain salt of n-butylthiomethyl-penicillin is prepared with procain hydrochloride, and after repeated recrystallization from methanol, prisms obtained and represent m. p. 110°C and potency 2,100 u/mg.

(5) The stabilities (half-life period) of n-butylthiomethyl-penicillin in aqueous solution at various pH, are determined and it is found that n-butylthiomethyl-penicillin is more stable than penicillin-G.

(6) The effective distribution coefficients of n-butylthiomethyl-penicillin between water and organic solvents at 0°C, pH 2.5 are determined. Butylacetate and ether give larger values and chloroform smaller value than that of penicillin-G.