Title	Strong shock waves in the electromagnetic shock tube
Sub Title	
Author	飯吉, 厚夫(liyoshi, Atsuo)
Publisher	慶応義塾大学藤原記念工学部
Publication year	1965
Jtitle	Proceedings of the Fujihara Memorial Faculty of Engineering Keio University (慶応義塾大学藤原記念工学部研究報告). Vol.18, No.71 (1965.),p.69(5)- 69(5)
JaLC DOI	
Abstract	
Notes	Summaries of Doctor and Master Theses Doctor of Engineering, 1965
Genre	Departmental Bulletin Paper
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=KO50001004-00180071- 0005

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Strong Shock Waves in the Electromagnetic Shock Tube

Atsuo IIYOSHI*

The strong shock waves such as produced in the electromagnetic shock tubes were studied experimentally and theoretically.

In the first part, the shock waves were observed with a rotaing camera, a photomultiprier and a piezo electric probe. Shock speeds were found to be in proportion to a factor $V/\sqrt{p_0}$ within the observed ranges of initial pressure (p_0) from 0.1 to 10 mmHg and of dischage voltage from 5 to 25 KV. This results agreed qualitatively with those of a simple analytical consideration based on a pinch expansion model. It was shown that behind a typical strong shock wave gas was partially dissociated and ionized, but it was practically observed as a local luminous region (radiation overshoot phenomenon). This radiation overshoot was also found to be mainly caused by radiation cooling.

In the second part, a unified theoretical analysis on the strong shock wave in Argon accompanying ionization phenomena were presented. In the relaxation phenomena accompanying dissociation and ionization, a considerable duration of time is required until the equilibrium is attained, and thus the shock front has certain thickness in strong contrast to the weak shock front which is taken mathematically as a discontinuous surface in the ordinary shock theory. The equilibrium state hehind the shock wave was examined numerically on the basis of Saha's equation and the Rankine-Hugoniot relations which was modified to ionized gases. Further, the analysis of the inner structure of the shock front were carried out taking account of the kinetics of the approach to equilibrium of ionization, which showed that atom temperature was higher than electron temperature in the relaxation region and the relaxation time was very sensitive to the electron temperature.

*飯 吉 厚 夫