

Title	Synthetic studies on aminosugar-glucosides and their derivatives : on the glucosides of deoxy-streptamine
Sub Title	
Author	土屋, 務(Tsuchiya, Tsutomu)
Publisher	慶應義塾大学藤原記念工学部
Publication year	1965
Jtitle	Proceedings of the Fujihara Memorial Faculty of Engineering Keio University (慶應義塾大学藤原記念工学部研究報告). Vol.18, No.71 (1965.) ,p.80(16)- 80(16)
JaLC DOI	
Abstract	
Notes	Summaries of Doctor and Master Theses Doctor of Engineering, 1965
Genre	Departmental Bulletin Paper
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=KO50001004-00180071-0016

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Synthetic Studies on Aminosugar-glucosides and Their Derivatives

—On the Glucosides of Deoxy-streptomine—

Tsutomu TSUCHIYA*

Recently, some antibiotics which are of practical importance have been found to be the glucosides containing amino-sugars, and this stimulated the chemists to make investigations on them.

This thesis concerns with the relationships between the structures of the amino-sugar-glucosides and their antibacterial activities as well as their syntheses.

(1) A partial degradation product of kanamycin, 4-0-(6-Amino-6-deoxy- α -D-glucosyl)-deoxystreptomine has been isolated and found to have antibacterial activity; this led to the conclusion that this part of kanamycin was the origin of the activity of kanamycin.

(2) A number of combinations of aminosugars and deoxystreptomine were subjected to the procedure so called "acid reversion", and it has been found that some new α -glucosides obtained by the method showed antibacterial activity.

(3) One of the above products which has strong antibacterial activity was isolated by means of paperchromatographic technique followed by some procedures and proved to be 0-(2-Amino-2-deoxy- α -D-glucosyl) deoxystreptomine.

(4) Kanamycin has seven alcoholic hydroxyl groups, one of which is primary and others are all secondary. The ease of arylsulfonation of this primary hydroxyl group has been utilized to the synthesis of deoxykanamycin and chlorodeoxykanamycin. Both the newly synthesized kanamycin derivatives have almost the same antibacterial activity as kanamycin and it has been found that the former has lower toxicity than kanamycin itself.

(5) Many derivatives of substituted or nonsubstituted kanamycin-N-methanesulfonates have been synthesized and the cause of the difference of their abilities of lowering the toxicity was discussed.

(6) Examination of the relationships between the conformations of the amino-sugar-glucosides and the antibacterial activities led to a conclusion that several special stereochemical conformations are essential to their antibacterial activities.

*土屋 務