| Title | Trapped electron velocity distribution function in plasmas : I. Positiveness of the distribution function in electron plasma |
|------------------|--|
| Sub Title | |
| Author | 小笠原, 正忠(Ogasawara, Masatada) 白井, 正司(Shirai, Shoji) |
| Publisher | 慶応義塾大学工学部 |
| Publication year | 1971 |
| Jtitle | Keio engineering reports Vol.24, No.7 (1971.) ,p.105- 108 |
| JaLC DOI | |
| Abstract | Sufficient condition for the positiveness of the trapped electron velocity distribution function in electron plasma is analyzed in the case of one-dimensional stationary wave solution of Vlasov and Poisson equations. If the electrostatic potential φ has a value between φ min and some critical value φ c, the sufficient condition for the positiveness is satisfied. The critical potential φ are evaluated in the limits of small and large φ . |
| Notes | |
| Genre | Departmental Bulletin Paper |
| URL | https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=KO50001004-00240007- 0105 |

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

KEIO ENGINEERING REPORTS VOL. 24 NO. 7

TRAPPED ELECTRON VELOCITY DISTRIBUTION FUNCTION IN PLASMAS I.

BY

MASATADA OGASAWARA and SHOJI SHIRAI

FACULTY OF ENGINEERING KEIO UNIVERSITY YOKOHAMA 1971

TRAPPED ELECTRON VELOCITY DISTRIBUTION FUNCTION IN PLASMAS

I. Positiveness of the Distribution Function in Electron Plasma

MASATADA OGASAWARA AND SHOJI SHIRAI

Dept. of Instrumentation Engineering, Keio University, Yokohama 223, Japan

(Received Dec. 22, 1971)

ABSTRACT

Sufficient condition for the positiveness of the trapped electron velocity distribution function in electron plasma is analyzed in the case of one-dimensional stationary wave solution of Vlasov and Poisson equations. If the electrostatic potential ϕ has a value between ϕ_{\min} and some critical value ϕ_c , the sufficient condition for the positiveness is satisfied. The critical potential ϕ_c are evaluated in the limits of small and large ϕ .

1. Introduction

The stationary one-dimensional nonlinear electrostatic wave has been studied by BERNSTEIN, GREENE and KRUSKAL (1957) with use of Vlasov and Poisson equations. The solutions of the basic equations were given as follows: If the potential and the distribution functions of ion and untrapped electron were given, Poisson equation allowed the distribution of the trapped electron to be calculated. The latter distribution function, however, is not always positive. An example was given by HATORI and SUGIHARA (1970). The trapped electron distribution was shown to be negative for some ranges of parameters of the untrapped electron velocity distribution.

SCHAMEL (1971) analyzed a necessary condition for the positiveness by assuming the trapped electron velocity distribution to be isotropic. Instead of the Poisson equation, TASSO (1969) used quasi-neutrality condition and showed that a sufficient condition for the positiveness was satisfied in an interval of potential. But his theory cannot take account of potential forms, i.e., sinusoidal, solitary and shocklike etc.. The aim of the present paper is to give a simple expression of a sufficient condition for the positiveness of the trapped electron velocity distribution function in an electron plasma.

2. Positiveness of $f_t(E)$

The basic equations for stationary electrostatic waves are

$$v \frac{\partial f(x,v)}{\partial x} + \frac{e}{m} \frac{d\phi(x)}{dx} \frac{\partial f(x,v)}{\partial v} = 0, \qquad (1)$$

$$\frac{d^2\phi(x)}{dx^2} = 4\pi e \int_{-\infty}^{\infty} dv f(x,v) - 4\pi e n_0, \qquad (2)$$

where f and ϕ are the electron velocity distribution function in the wave frame and potential, respectively. Ion density is denoted by n_0 . The general solution of equation (1) is

$$f = f(E), \qquad E = \frac{1}{2} mv^2 - e\phi.$$
 (3)

Substituting (3) into (2) and denoting "trapped" and "untrapped" by subscripts t and u, we obtain

$$\int_{-e\phi}^{-e\phi\min} dE f_t(E) [2m(E+e\phi)]^{-\frac{1}{2}} = g(e\phi), \qquad (4)$$

where

$$g(e\phi) = -\frac{N(e\phi)}{4\pi e} + n_0 - \int_{-e\phi_{\min}}^{\infty} dE f_u(E) [2m(E+e\phi)]^{-\frac{1}{2}}$$
(5)

is the density of trapped electron at x corresponding to $\phi(x)$. Because of (4), $g(e\phi_{\min})=0$. If the charge density $N(e\phi)=-d^2\phi/dx^2$ and n_0 and $f_u(E)$ are prescribed, $g(e\phi)$ is a known function and (4) is an integral equation of the convolution type for $f_t(E)$ and following BERNSTEIN, GREENE and KRUSKAL (1957), the solution is given by way of Laplace transform

$$f_{\ell}(E) = \frac{\sqrt{2m}}{\pi} \int_{-e\phi_{\min}}^{-E} dV \frac{dg(V)}{dV} \frac{1}{\sqrt{-E-V}}, \quad E < -e\phi_{\min}.$$
(6)

From the expression (6), it follows that $\frac{dg}{dV} > 0$ is a sufficient condition for the positiveness of $f_t(E)$. From (5) we have

$$\frac{dg(V)}{dV} = -\frac{1}{4\pi e} \frac{dN(V)}{dV} + \frac{1}{2\sqrt{2m}} \int_{-e\phi_{\min}}^{\infty} dE \frac{f_u(E)}{(E+V)^{\frac{3}{2}}}.$$
 (7)

Then the sufficient condition for the positiveness is given by

Trapped Electron Velocity Distribution Function in Plasmas I.

$$\frac{dg}{dV} \ge \frac{1}{2\sqrt{2m}} \int_{-e\phi_{\min}}^{\infty} dE \frac{f_u(E)}{(E+V)^{\frac{1}{2}}} - \frac{M}{4\pi e} > 0, \qquad (8)$$

where

$$M = Maximum \ of \ \frac{dN}{dV} \,. \tag{9}$$

For physically reasonable potential ϕ , M is positive. [c.f. VALEO, OBERMAN and KRUS-KAL (1969).] If $f_u(E)$ is defined in the whole range of E, the first term of (8) goes to ∞ as $V \rightarrow e\phi_{\min}$ and decreases monotonically to zero as $V \rightarrow \infty$. Then Vhas a critical value $V_c(=e\phi_c)$ beyond which $\frac{dg(V)}{dV} < 0$. The result is similar to that of TASSO (1969).

In the limit of small and large ϕ , the critical potential ϕ_c can be derived analytically as follows: We have small ϕ expansion by making partial integration in (8),

$$\frac{1}{\sqrt{2m}} \left\{ \frac{f_u(-e\phi_{\min})}{\sqrt{e(\phi-\phi_{\min})}} + 2\sqrt{e(\phi-\phi_{\min})} f'_u(-e\phi_{\min}) + \cdots \right\} - \frac{M}{4\pi e} > 0.$$
(10)

This expansion has meaning if $e(\phi - \phi_{\min}) \ll V_T$, V_T being the width of $f_u(V)$. With use of (10), the critical ϕ_c in the small ϕ limit is given by

$$\phi_c - \phi_{\min} = \frac{1}{e} \left[\frac{4\pi e f_u(-e\phi_{\min})}{\sqrt{2m} M} \right]^2.$$
(11)

When V is much larger than $V_0 + V_T$, V_0 being defined by $f_u(V_0) = \text{maximum}$, it follows that

$$\frac{1}{\sqrt{2m}} \int_{-e\phi_{\min}}^{\infty} \frac{f_u(E)dE}{(E+V)^{\frac{3}{2}}} \approx \frac{n_u}{V},$$
(12)

where n_u , the untrapped electron number density, is defined by

$$n_u = \int_{-e\phi_{\min}}^{\infty} \frac{f_u(E)dE}{\sqrt{2m(E+V)}} \,. \tag{13}$$

According to MONTGOMERY and JOYCE (1969), n_u are connected with n_0 , unperturbed number density, by

$$n_u = \frac{n_0}{\sqrt{1 + \frac{V}{V_T}}} \approx n_0 \sqrt{\frac{V_T}{V}} \,. \tag{14}$$

Then in the limit of large ϕ , it follows from (8), (12) and (14) that

$$\phi_c = \frac{1}{e} \left[\frac{2\pi e n_0 \sqrt{V_T}}{M} \right]^{\frac{2}{3}}.$$
(15)

For the intermediate value of ϕ , the following expression for the positiveness criterion will be convenient

$$\frac{1}{\sqrt{2me(\phi-\phi_{\min})}} \frac{f_u(-e\phi_{\min})+4f_u(3e\phi-4e\phi_{\min})}{6} - \frac{M}{4\pi e} > 0, \qquad (16)$$

where Simpson's formula has been used for evaluating the integral.

For examples we take a potential given by

$$\phi(x) = \frac{\phi_0}{\cosh^l(kx)}.$$
(17)

This expresses a solitary wave if k is real and l is a positive integer. It follows from (17) that

$$M = \frac{k^2}{e} (3l+2), \qquad \phi_{\min} = 0.$$
(18)

When k is pure imaginary and l=-1, the potential $\phi(x)$ given by (17) reduces to a sinusoidal one, and it follows that

$$M = \frac{k^2}{e}, \qquad \phi_{\min} = -\phi_0.$$
 (19)

It is to be noted that in both cases M are positive and the critical values ϕ_c exist.

REFERENCES

BERNSTEIN I. B., GREENE J. M. and KRUSKAL M. D. (1957): Exact Nonlinear Plasma Oscillations, Phys. Rev. 108, 546.

HATORI T. and SUGIHARA R. (1970): Private Communication.

- MONTGOMERY D. and JOYCE G. (1970): Shock-Like Solutions of Electrostatic Vlasov Equation, J. Plasma Physics 3, 1.
- SCHAMEL H. (1971): Stationary Solutions of the Electrostatic Vlasov Equation, Plasma Physics 13, 491.

TASSO H. (1969): Nonlinear Quasi-Neutral Electrostatic Plasma Waves and Shock Waves, Plasma Physics 11, 663.

VALEO E. J., OBERMAN C. and KRUSKAL M. D. (1969): Property of Large-Amplitude Steady Electrostatic Waves, Phys. Fluids 12, 1246.