Title	On the strong law of large numbers for harmonizable stochastic processes
Sub Title	
Author	有本, 彰雄(Arimoto, Akio)
Publisher	慶応義塾大学工学部
Publication year	1972
Jtitle	Keio engineering reports Vol.25, No.8 (1972.),p.101-111
JaLC DOI	
Abstract	The harmonizable stochastic processes $X(t) = \blacksquare Z(dx)$ are additively built up by elementary harmonic oscilations $\blacksquare Z(dx)$ , each of which has an angular frequency x, while the random amplitude and the random phase are determined by the elementary increment $Z(dx)$ . It is well known that the necessary and sufficient condition for the validity of the weak law of large numbers for such a process is described in terms of the spectral distribution $EZ(dx)$ . However, for the strong law of large numbers, sufficient conditions have been given in terms of covariance functions for weakly stationary processes or some non-stationary stochastic processes. In this paper a sufficient condition for the strong law of large numbers is given in terms of the spectral distribution for harmonizable stochastic processes.
Notes	
Genre	Departmental Bulletin Paper
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=KO50001004-00250008- 0101

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

## KEIO ENGINEERING REPORTS VOL. 25 NO. 8

# ON THE STRONG LAW OF LARGE NUMBERS FOR HARMONIZABLE STOCHASTIC PROCESSES

BY

AKIO ARIMOTO

FACULTY OF ENGINEERING KEIO UNIVERSITY YOKOHAMA 1972

### ON THE STRONG LAW OF LARGE NUMBERS FOR HARMONIZABLE STOCHASTIC PROCESSES

Ακίο Αγιμότο

Dept. of Administration Engineering, Keio University, Yokohama 223, Japan\*

(Received Oct. 5, 1972)

#### ABSTRACT

The harmonizable stochastic processes  $X(t) = \int_{-\infty}^{\infty} e^{itx}Z(dx)$  are additively built up by elementary harmonic oscilations  $e^{itx}Z(dx)$ , each of which has an angular frequency x, while the random amplitude and the random phase are determined by the elementary increment Z(dx). It is well known that the necessary and sufficient condition for the validity of the weak law of large numbers for such a process is described in terms of the spectral distribution  $EZ(dx)\overline{Z(dy)}$ . However, for the strong law of large numbers, sufficient conditions have been given in terms of covariance functions for weakly stationary processes or some non-stationary stochastic processes. In this paper a sufficient condition for the strong law of large numbers is given in terms of the spectral distribution for harmonizable stochastic processes.

1. Let

(1.1) 
$$X(t) = \int_{-\infty}^{\infty} e^{itx} Z(dx)$$

be a harmonizable stochastic process, in which Z(S) is an additive stochastic set function defined for each Borel set S on the real axis such that

$$(1.2) EZ(S)=0,$$

(1.3) 
$$EZ(S_1)\overline{Z(S_2)} = \int_{x \in S_1} \int_{y \in S_2} dF(x, y),$$

where F(x, y) is a function of limited total variation over the whole plane, that is,

(1.4) 
$$\int_{-\infty}^{\infty} |dF(x,y)| < \infty.$$

<sup>\*</sup> Now at Musashi Institute of Technology, Dept. of Industrial Engineering.

#### Ακίο Αγιμότο

X(t) is a process of the second order with EX(t)=0 and the covariance function of X(t) is given by

(1.5) 
$$r(s,t) = EX(s)\overline{X(t)}$$
$$= \int_{-\infty}^{\infty} e^{i(sx-ty)} dF(x,y),$$

which is continuous on  $-\infty < s, t < \infty$  and F(x, y) is hermitian symmetric;  $F(x, y) = \overline{F(y, x)}$ . F(x, y) is called the spectral distribution function of X(t). Such a process was first introduced by LOÈVE (1948) and some applications of this process were discussed in BLANC-LAPPIERRE and FORTET (1953). In LOÈVE (1948), he has shown that if X(t) is a harmonizable stochastic process, then

(1.6) 
$$Y(u) = \lim_{T \to \infty} \frac{1}{2T} \int_{-T}^{T} e^{-iut} X(t) dt$$

exists for every u, EY(u) = 0 and

$$(1.7) EY(u) \overline{Y(v)}$$

$$=F(u+0, v+0)-F(u-0, v+0)-F(u+0, v-0)+F(u-0, v-0).$$

That is, the weak law of large numbers for X(t) always holds, whenever F(x, y) is continuous at origin.

Now we talk about the strong law of large numbers. Loève (1948) and CRAMÉR-LEADBETTER (1967) gave the sufficient conditions for the validity of the strong law of large numbers for general stochastic processes of second order.

THEOREM A. (LOÈVE (948)) Suppose that X(t) is a stochastic process which is continuous in the mean over all the finite segment. If, either

(I) for sufficiently large t,

$$(1.8) r(t,t) < C < \infty and$$

(1.9) 
$$\frac{1}{T^{2-c}}\int^{T}\int^{T}r(u,v) \, du dv < D < \infty,$$

where C>0, c>0, D are some constants, or

(II) with probability one,

$$(1.10) |X(t)| < C' < \infty, -\infty < t < \infty and$$

(1.11) 
$$\int_{-\infty}^{\infty} \frac{dT}{T^3} \int_{-\infty}^{T} |r(u,v)| \, du \, dv < D' < \infty,$$

where C'>0, D'>0 are some constants, then we have

(1.12) 
$$\lim_{T \to \infty} \frac{1}{T} \int^T X(t) dt = 0$$

with probability one.

THEOREM B. (CRAMÉR-LEADBETTER (1967)) Suppose that r(s, t) is continuous for  $s, t \ge 0$  and that

(1.13) 
$$|r(s,t)| < K \frac{s^{\alpha} + t^{\alpha}}{1 + |s-t|^{\beta}},$$

where K,  $\alpha$  and  $\beta$  are constants such that  $K>0, 0\leq 2\alpha < \beta < 1$ . Then we have

(1.14) 
$$\lim_{T \to \infty} \frac{1}{T} \int_0^T X(t) dt = 0$$

with probability one.

These can be applied to the case of harmonizable processes. On the other hand VERBITSKAYA (1964) gave the condition for the strong law of large numbers for weakly stationary processes which were better than the above LOÈVE and CRAMÉR-LEADBETTE's conditions applied to weakly stationary processes. However, a remarkable fact is that the method of the proof of VERBITSKAYA could be applied to the more general processes not necessarily stationary and in fact, KAWATA (1972) has noticed the following result applying it to some other class of non-stationary processes:

THEOREM C. Let X(t) be a stochastic process of the second order with  $EX(t) = 0, -\infty < t < \infty$ . If

(I)

(1.15) 
$$\int_{n}^{n+1} E|X(t)|^{2} dt < K,$$

for all integer n with  $-\infty < n < \infty$ , K being a constant independent of n, and

(II) there is a non-negative function  $g(u), 0 \leq u < \infty$ , such that

(1.16) 
$$\left|\int_{0}^{x}\int_{0}^{x}r(u+\tau,v+\tau)\,dudv\right| \leq \int_{0}^{x}g(u)du$$

for every  $-\infty < \tau < \infty, 0 \leq x < \infty$ , where

(1.17) 
$$\int_{1}^{\infty} \frac{\log^2 x}{x^2} g(x) dx < \infty.$$

Then

(1.18) 
$$\lim_{T \to \infty} \frac{1}{T} \int_0^T X(t) dt = 0$$

#### with probability one.

In this paper, we shall, following KAWATA's line, handle the problem and give sufficient conditions in terms of the spectral distribution F(x, y) for the strong law of large numbers in the case of harmonizable processes.

#### Ακίο Αγιμότο

2. Let X(t) be a given harmonizable process with  $EX(t)=0, -\infty < t < \infty$ , and the spectral distribution function F(x, y). We begin with the following lemma.

LEMMA 1. Suppose that for  $u \rightarrow 0+$ ,

(2.1) 
$$\int_{D_1} \int \frac{1}{|x|^a |y|^b} |dF(x, y)| = O\left(\frac{1}{u^{a+b} |\log u|^{3+\delta}}\right),$$

(2.2) 
$$\int_{D_2} \int \frac{1}{|x|^c |y|^d} |dF(x, y)| = O\left(\frac{1}{u^{c+d} |\log u|^{3+\delta}}\right),$$

(2.3) 
$$\int_{D_3} \int \frac{1}{|x|^e |y|^f} |dF(x, y)| = O\left(\frac{1}{u^{e+f} |\log u|^{3+\delta}}\right),$$

for some  $a, b, c \in [0, \infty)$  and some  $d, e, f \in [0, 1]$  and some  $\delta > 0$ , where

$$D_1 = \{(x, y): |x| \le u, |y| \le u\},$$
  
 $D_2 = \{(x, y): |x| \le u, u < |y|\}$ 

and

$$D_3 = \{(x, y): u < |x| \le 1, u < |y| \le 1\}.$$

Then

(2.4) 
$$\int_1^\infty \frac{\log^2 t}{t^3} h(t,\tau) dt < \infty,$$

where

(2.5) 
$$h(t,\tau) = \max_{0 \le s \le t} \left\{ \int_0^s \int_0^s r(u+\tau,v+\tau) \, du \, dv \right\}, -\infty < \tau < \infty.$$

Proof.

We can easily see by (1.4) and (1.5) that

(2.6) 
$$\int_{0}^{t} \int_{0}^{t} r(u+\tau, v+\tau) du dv$$
$$= 4 \iint_{-\infty}^{\infty} e^{i\left(\tau+\frac{t}{2}\right)(x-y)} \frac{\sin \frac{xt}{2} \sin \frac{yt}{2}}{xy} dF(x, y)$$

Therefore, we have

(2.7) 
$$\left| \int_{0}^{t} \int_{0}^{t} r(u+\tau, v+\tau) \, du dv \right|$$
$$\leq 4 \int \int_{-\infty}^{\infty} \left| \frac{\sin \frac{xt}{2}}{x} \frac{\sin \frac{yt}{2}}{y} \right| |dF(x, y)|$$

which is split into five parts for large t > 0

$$\int_{D_1^*} \int + \int_{D_2^*} \int + \int_{D_3^*} \int + \int_{D_4^*} \int + \int_{D_5^*} \int ,$$

where

$$D_{1}^{*} = \left\{ (x, y) \colon |x| \leq \frac{2}{t}, |y| \leq \frac{2}{t} \right\},$$
$$D_{2}^{*} = \left\{ (x, y) \colon |x| \leq \frac{2}{t}, |y| > \frac{2}{t} \right\},$$
$$D_{3}^{*} = \left\{ (x, y) \colon |x| > \frac{2}{t}, |y| \leq \frac{2}{t} \right\},$$
$$D_{4}^{*} = \left\{ (x, y) \colon \frac{2}{t} < |x| \leq 1, \frac{2}{t} < |y| \leq 1 \right\},$$

and

$$D_5^* = (-\infty, \infty) \times (-\infty, \infty) - D_1^* \cup D_2^* \cup D_3^* \cup D_4^*.$$

Since  $|\sin x| \leq |x|$ , we have

(2.8) 
$$\int_{0}^{2/t} \int_{0}^{2/t} \frac{\sin \frac{xt}{2} \sin \frac{yt}{2}}{xy} |dF(x,y)| \leq \left(\frac{t}{2}\right)^{2-a-b} \int_{0}^{2/t} \int_{0}^{2/t} \frac{1}{x^{a}y^{b}} |dF(x,y)|,$$

(2.9) 
$$\int_{0}^{2/t} \int_{2/t}^{\infty} \frac{\sin \frac{xt}{2} \sin \frac{yt}{2}}{xy} |dF(x,y)| \leq \left(\frac{t}{2}\right)^{1-c} \int_{0}^{2/t} \int_{2/t}^{\infty} \frac{1}{x^{c}y} |dF(x,y)|$$

which is, since

$$\frac{1}{y} = \frac{1}{y^d y^{1-d}} \le \frac{1}{y^d} \left(\frac{t}{2}\right)^{1-d} \quad \text{for} \quad y \ge \frac{2}{t},$$
$$\le \left(\frac{t}{2}\right)^{2-c-d} \int_0^{2/t} \int_{2/t}^\infty \frac{1}{x^c y^d} |dF(x,y)|,$$
$$\int_{0}^1 \int_{0}^1 \frac{\sin \frac{xt}{2} \sin \frac{yt}{2}}{xy} |dF(x,y)| \le \int_{0}^1 \int_{0}^1 \frac{1}{xy} |dF(x,y)|$$

$$\leq \left(\frac{t}{2}\right)^{2-e-f} \int_{2/t}^{1} \int_{2/t}^{1} \frac{1}{x^{e}y^{f}} |dF(x,y)|$$

and

(2.10)

(2.11) 
$$\int_{1}^{\infty} \int_{1}^{\infty} \frac{\sin \frac{xt}{2} \sin \frac{yt}{2}}{xy} |dF(x,y)| \leq \int_{0}^{\infty} \int_{0}^{\infty} |dF(x,y)|.$$

We have similar estimates of the integrals in other quadrants. Thus, we have

(2.12) 
$$\left| \frac{1}{4} \int_{0}^{t} \int_{0}^{t} r(u+\tau, v+\tau) \, du dv \right|$$

$$\leq \left( \frac{t}{2} \right)^{2-a-b} \int_{D_{1}^{*}} \int_{1}^{1} \frac{1}{|x|^{a} |y|^{b}} |dF(x, y)| + 2 \left( \frac{t}{2} \right)^{2-c-d} \int_{D_{2}^{*}} \int_{1}^{1} \frac{1}{|x|^{c} |y|^{d}} |dF(x, y)|$$

Ακιο Αγιμότο

$$+\left(\frac{t}{2}\right)^{2-e-f}\int_{D_{4}^{*}}\int_{|x|^{e}|y|^{f}}|dF(x,y)|+\int_{-\infty}^{\infty}|dF(x,y)|.$$

Now we can easily see

$$\left|\int_0^t \int_0^t r(u+\tau, v+\tau) \, du dv\right| \leq t^2 \int \int_{-\infty}^\infty |dF(x, y)|.$$

Hence, taking u=2/t in (2.1), (2.2) and (2.3), we have, for large t,

(2.13) 
$$h(t,\tau) \leq C \frac{t^2}{\log^{3+\delta} t}$$

for some constant C>0, which completes the proof.

Here we shall give the conditions for the strong law of large numbers for X(t) in the following theorem.

THEOREM 1. Suppose that the spectral distribution function F(x, y) of a process X(t) satisfies the following conditions: for  $\delta > 0$ ,

(2.14) 
$$\int_{|x| \le u} \int_{|y| \le u} |dF(x, y)| = O\left(\frac{1}{|\log u|^{3+\delta}}\right)$$

(2.15) 
$$\int_{|x| \le u} \int_{|y| > u} \frac{1}{|y|} |dF(x, y)| = O\left(\frac{1}{|u| \log |u|^{3+\delta}}\right),$$

and

(2.16) 
$$\int_{|x|>u} \int_{|y|>u} \frac{1}{|xy|} |dF(x, y)| = O\left(\frac{1}{u^2 |\log u|^{3+\delta}}\right)$$

for small u>0. Then

(2.17) 
$$\lim_{T \to \infty} \frac{1}{2T} \int_{-T}^{T} X(t) dt = 0$$

with probabilility one.

*Proof.* To prove the theorem we use the method of VERBITSKAYA (1964). We shall prove only that  $\frac{1}{T} \int_0^T X(t) dt \to 0$ , since  $\frac{1}{T} \int_{-T}^0 X(t) dt \to 0$  can be proved in the same manner. Let

(2.18) 
$$S(T) = \int_0^T X(t) dt, T \ge 1.$$

For any given T, there exists a positive integer n such that  $n \le T \le n+1$ . Then, we have

(2.19) 
$$\left|\frac{1}{T}\int_{0}^{T}X(t)dt\right| \leq \left|\frac{1}{n}\int_{0}^{n}X(t)dt\right| + \frac{1}{n}\int_{n}^{n+1}|X(t)|dt.$$

The second term of the right hand side approaches zero with probability one when n tends to infinity, because

(2.20) 
$$\sum_{n} E\left(\frac{1}{n}\int_{n}^{n+1}|X(t)|dt\right)^{2}$$
$$\leq \sum_{n} \frac{1}{n^{2}} E\int_{n}^{n+1}|X(t)|^{2}dt$$
$$= \sum_{n} \frac{1}{n^{2}} E\int_{-\infty}^{\infty}|dF(x,y)| < \infty.$$

Hence, it suffices to show that

(2.21) 
$$\lim_{n \to \infty} \frac{S(n)}{n} = 0$$

with probability one.

Now any positive integer n can be uniquely represented in the form

$$(2.22) n=2^{k_1}+2^{k_2}+2^{k_3}+\cdots+2^{k_s}, k_1>k_2>k_3\cdots>k_s\geq 0,$$

 $k'_i$ s being integers. Then we can write

$$(2.23) |S(n)| \leq |S(2^{k_1})| + |S(2^{k_1} + 2^{k_2}) - S(2^{k_1})| + \dots + |S(2^{k_1} + (2^{k_2 - k_3} + 2^{k_3 - k_3} + \dots + 2^{k_{3-1} - k_3} + 1)2^{k_3}) - S\{2^{k_1} + (2^{k_2 - k_3} + \dots + 2^{k_{3-1} - k_3})2^{k_3}\}|.$$

Using this relation, we have only to show that the series, for any  $\varepsilon > 0$  fixed,

(2.24) 
$$\sum_{k=1}^{\infty} P\left\{ \left| \frac{1}{2^k} S(2^k) \right| \ge \frac{\varepsilon}{k+1} \right\},$$

(2.25) 
$$\sum_{k=1}^{\infty} \sum_{p=0}^{k-1} \sum_{q=0}^{2^{k-p-1}} P\left\{ \left| \frac{1}{2^k} (S(2^k + (q+1)2^p) - S(2^k + q2^p)) \right| \ge \frac{\varepsilon}{k+1} \right\}$$

are convergent (VERBITSKAYA 1964). But by the Tchebychev's inequality, we have

(2.26) 
$$\sum_{k=1}^{\infty} P\left\{ \left| \frac{S(2^k)}{2^k} \right| \ge \frac{\varepsilon}{k+1} \right\}$$

$$\leq \frac{1}{\varepsilon^2} \sum_{k=0}^{\infty} \frac{(k+1)^2}{2^{2k}} \int_0^{2^k} \int_0^{2^k} r(u,v) \, du \, dv$$

which is

$$\leq \frac{1}{\varepsilon^2} \sum_{k=1}^{\infty} \frac{(k+1)^2}{2^{2k}} h(2^k, 0)$$
$$\leq \frac{C}{\varepsilon^2} \int_1^\infty \frac{\log^2 x}{x^3} h(x, 0) dx,$$

where  $h(x, \tau)$  is the one in (2.5) and C is a positive constant. Here the last term

converges by virtue of Lemma 1. For (2.25) we have, again using the Tchebychev's inequality,

(2.27) 
$$\sum_{k=1}^{\infty} \sum_{p=0}^{k-1} \sum_{q=0}^{2^{k-p-1}} P\left\{ \left| \frac{1}{2^{k}} (S(2^{k} + (q+1)2^{p}) - S(2^{k} + q2^{p})) \right| \ge \frac{\varepsilon}{k+1} \right\}$$
$$\le \frac{1}{\varepsilon^{2}} \sum_{k=1}^{\infty} \sum_{p=0}^{k-1} \sum_{q=0}^{2^{k-p-1}} \frac{(k+1)^{2}}{2^{2^{k}}} \left| \int_{0}^{2^{p}} \int_{0}^{2^{p}} r(2^{k} + q2^{p} + u, 2^{k} + q2^{p} + v) du dv$$

which is

 $\leq rac{C}{arepsilon^2} \sum_{k=1}^{\infty} \sum_{p=0}^{k-1} rac{(k+1)^2}{2^{k+p}} \ h(2^p, au_0),$ 

where  $\tau_0$  is such that

(2.28) 
$$h(2^{p}, \tau_{0}) = \max_{0 \le q \le 2^{\kappa-p}-1} \{|h(2^{p}, 2^{k}+q2^{p})|\},$$

which is

$$\leq \frac{C}{\varepsilon^2} \sum_{k=1}^{\infty} \frac{(k+1)^2}{2^{2k}} h(2^k, \tau_0)$$
$$\leq \frac{C}{\varepsilon^2} \int_1^{\infty} \frac{\log^2 x}{x^3} h(x, \tau_0) dx$$

for some constant C and the last term is finite because of Lemma 1. We have now completed the proof.

Each of the conditions  $(2.14)\sim(2.16)$  can be replaced respectively by a little better one, each of  $(2.1)\sim(2.3)$ .

THEOREM 2. Suppose that the spectral distribution function F(x, y) satisfies the conditions  $(2, 1) \sim (2, 3)$ . Then

(2.29) 
$$\lim_{T \to \infty} \frac{1}{2T} \int_{-T}^{T} X(t) dt = 0$$

with probability one.

The proof is carried out in just the same way as that of Theorem 1. Let us now turn to the special case of a uniformly bounded process.

THEOREM 3. Suppose that a constant K exists such that with probability one,

$$(2.30) |X(t)| \leq K$$

holds for any t. If for  $u \rightarrow +0$ ,

(2.31) 
$$\int_{|x| \le u} \int_{|y| \le u} |dF(x, y)| = O\left(\frac{1}{|\log u|^{1+\delta}}\right),$$

(2.32) 
$$\int_{|x| \le u} \int_{|y| > u} \frac{1}{|y|} |dF(x, y)| = O\left(\frac{1}{u |\log u|^{1+\delta}}\right)$$

and

(2.33) 
$$\int_{|x|>u} \int_{|y|>u} \frac{1}{|xy|} |dF(x, y)| = O\left(\frac{1}{u^2 |\log u|^{1+\delta}}\right)$$

for  $\delta > 0$ , then we have

(2.34)

$$\lim_{T\to\infty}\frac{1}{2T}\int_{-T}^{T}X(t)dt=0$$

with probability one.

 $(2.31)\sim(2.33)$  can be replaced by a little better ones as in Theorem 2.

**Proof of Theorem 3.** The idea of the following proof is due to VERBITSKAYA. Partition the infinite interval  $[1, \infty)$  into subintervals  $[2^n, 2^{n+1}]$ ,  $n=0, 1, 2, \cdots$  and let  $\varepsilon$  be a small positive real number prescribed in an arbitrary manner. Choose an integral l such that  $K/l < \varepsilon/2$ . Now partition each interval  $[2^n, 2^{n+1}]$  into l equal parts. Number all the resulting division points according to the order of increasing magnitude. This yields a sequence  $\{T_n\}$  of division points,  $T_n \to \infty$  as  $n \to \infty$ .

Now suppose T is arbitrarily large. Then for any T>0, an integer n exists for which  $T_n \leq T \leq T_{n+1}$ . Let the interval  $[2^m, 2^{m+1}]$  contain both  $T_n$  and  $T_{n+1}$ . Then clearly  $T_{n+1}-T_n=2^m/l$ . Thus, for any  $\varepsilon>0$ , we have, by (2.30)

(2.35) 
$$\frac{1}{T_n} \int_{T_n}^{T_{n+1}} |X(t)| dt < \frac{\varepsilon}{2}.$$

Hence, we have only to show

(2.36) 
$$\sum_{n=1}^{\infty} P\left\{\frac{1}{T_n}\left|\int_0^{T_n} X(t)dt\right| \ge \frac{\varepsilon}{2}\right\} < \infty.$$

we can see that

(2.37) 
$$\sum_{n=1}^{\infty} P\left\{\frac{1}{T_n} \left| \int_0^{T_n} X(t) dt \right| \ge \frac{\varepsilon}{2} \right\}$$

$$= \sum_{j=0}^{l-1} \sum_{n=1}^{\infty} P\left\{\frac{1}{2^n + j\frac{2^n}{l}} \int_0^{2^n + j(2^n/l)} X(t) dt \right| \ge \frac{\varepsilon}{2}\right\}$$

and

(2.38) 
$$P\left\{\frac{1}{2^n+j\frac{2^n}{l}}\left|\int_0^{2^n+j(2^n/l)}X(t)dt\right| \ge \frac{\varepsilon}{2}\right\}$$

$$\leq \frac{2^2}{\varepsilon^2 \left(2^n + j\frac{2^n}{l}\right)^2} E \left| \int_0^{2^n + j(2^n/l)} X(t) dt \right|^2$$

$$\leq \frac{\left(2^{n}+j\frac{2^{n}}{l}\right)2^{2}}{\varepsilon^{2}\left(2^{n}+j\frac{2^{n}}{l}\right)^{3}}h(2^{n}+j2^{n}/l,0) \\ \leq \frac{4}{\varepsilon^{2}}\int_{2^{n+1+j(2^{n+1}/l)}}^{2^{n+1+j(2^{n+1}/l)}}\frac{1}{x^{3}}h(x,0)dx.$$

Therefore, the (2.36) holds if

(2.39) 
$$\int_{1}^{\infty} \frac{1}{t^3} h(t, 0) dt < \infty$$

and this is easily seen to be implied by  $(2.31)\sim(2.33)$ , which completes the proof.

Finally we give some remarks. Although our theorem is not really a generalization of Loève's theorem, the following consideration will tell us some connection between them.

For any harmonizable process

$$|\mathbf{r}(t,t)| \leq \int_{-\infty}^{\infty} |dF(x,y)|, \ -\infty < t < \infty,$$

holds and the conditions (2.1), (2.2) and (2.3) imply, by (2.13),

(2.41) 
$$\int_{0}^{t} \int_{0}^{t} r(u, v) du dv = O\left(\frac{t^{2}}{\log^{3+\delta} t}\right),$$

which is a condition weaker than (1.9). Furthermore, if the right hand sides of (2.1), (2.2) and (2.3) are replaced by the respective weaker conditions

$$O\left(\frac{1}{|u^{a+b}|\log u|^{1+\delta}}\right), O\left(\frac{1}{|u^{c+d}|\log u|^{1+\delta}}\right) \text{ and } O\left(\frac{1}{|u^{e+f}|\log u|^{1+\delta}}\right), \text{ then}$$

$$(2.42) \qquad \qquad \int_{0}^{t} \int_{0}^{t} r(u,v) \, du \, dv = O\left(\frac{t^2}{\log^{1+\delta} t}\right)$$

holds and hence

(2.43) 
$$\int_1^\infty \frac{dt}{t^3} \int_0^t \int_0^t r(u,v) du dv < \infty$$

holds. The condition (2.43) is obviously weaker than (1.11).

#### Acknowledgement

The author would like to express his thanks to Professor Tatsuo KAWATA of Keio University for his constant encouragement and many helpful suggestions.

#### REFERENCES

- BLANC-LAPPIERRE, A., and FORTET, R. (1953): Theorie des fonction aleatoires, Libraires de L'Academie de Medecine, Paris.
- CRAMÉR, H., and LEADBETTER, M. R. (1967): Stationary and related stochastic processes, Wiley, New York.
- KAWATA, T. (1972): On a class of linear processes, Read at the Second Japan-USSR Symposium (Kyoto), to appear.

- Loève, M. (1948): Fonctions aleatoires du second ordre, appendix to P. Levy, Processus stochastiques et movemont brownien, Gauthier Villars, Paris.
- VERBITSKAYA, I. N. (1964): On conditions for the strong law of large numbers to be applicable to second order stationary processes, *Theoriya Veroyatonostei i ee Primeneniya* 9 pp. 325-331.