

Title	A parametrization of Benders decomposition
Sub Title	
Author	Tone, Kaoru
Publisher	慶應義塾大学工学部
Publication year	1976
Jtitle	Keio engineering reports Vol.29, No.5 (1976. 6) ,p.53- 60
JaLC DOI	
Abstract	Benders decomposition is a multi-step procedure effective for solving mixed variables programming problems. In this paper, we shall present a parametrization of Benders decomposition with respect to a parameter on the right hand side of constraints.
Notes	
Genre	Departmental Bulletin Paper
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=KO50001004-00290005-0053

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

A PARAMETRIZATION OF BENDERS DECOMPOSITION

KAORU TONE

Dept. of Mathematics, Keio University, Yokohama, 223, Japan

(Received May 17, 1976)

ABSTRACT

Benders decomposition is a multi-step procedure effective for solving mixed variables programming problems.

In this paper, we shall present a parametrization of Benders decomposition with respect to a parameter on the right hand side of constraints.

1. Preliminaries —Basic Theorems of Benders Decomposition and Parametric Analysis by Primal-Dual Algorithm—

In our analysis, we use two procedures, as subroutines, namely Benders decomposition and a parametric analysis of linear programmings by primal-dual algorithm, which are summarized as follows.

1.1 Basic Theorems of Benders Decomposition

BENDERS (1962) has presented a partitioning procedure for solving mixed variables programming problems of the type

$$(1.1) \quad \text{Max } \{c'x + f(y) \mid Ax + F(y) \leq b, 0 \leq x \in R^p, y \in S\},$$

where $x \in R^p$ (the p -dimensional Euclidean space), $y \in R^q$ and S is an arbitrary subset of R^q . Furthermore, A is an (m, p) matrix, $f(y)$ is a scalar function and $F(y)$ an m -component vector function both defined on S , b and c are fixed vectors in R^m and R^p , respectively, and prime denotes transposition.

His basic idea is a partitioning of the given problem into two subproblems: a programming problem (which may be linear, nonlinear, discrete, etc.) defined on S , and a linear programming problem defined on R^p .

In this connection, he defined two sets C and G as follows:

(a) A polyhedral convex cone C in R^{m+1} :

$$(1.2) \quad C = \{(u_0, u) | A'u - cu_0 \geq 0, u \geq 0, u_0 \geq 0\}.$$

(b) A set G in R^{n+1} :

$$(1.3) \quad G = \bigcap_{(u_0, u) \in C} \{(x_0, y) | u_0 x_0 + u'F(y) - u_0 f(y) \leq u'b, y \in S\}.$$

Then, he states the basic theorem for a partitioning procedure:

Theorem 1.1

(1) Problem (1.1) is not feasible if and only if the programming problem

$$(1.4) \quad \text{Max}\{x_0 | (x_0, y) \in G\}$$

is not feasible, i.e. if and only if the set G is empty.

(2) Problem (1.1) is feasible without having an optimum solution, if and only if Problem (1.4) is feasible without having an optimum solution.

(3) If (\bar{x}, \bar{y}) is an optimum solution of Problem (1.1) and

$$\bar{x}_0 = c'\bar{x} + f(\bar{y}),$$

then (\bar{x}_0, \bar{y}) is an optimum solution of Problem (1.4) and \bar{x} is an optimum solution of the linear programming problem

$$(1.5) \quad \text{Max}\{c'x | Ax \leq b - F(\bar{y}), x \geq 0\}.$$

(4) If (\bar{x}_0, \bar{y}) is an optimum solution of Problem (1.4), then Problem (1.5) is feasible and the optimum value of the objective function in this problem is equal to $\bar{x}_0 - f(\bar{y})$. If \bar{x} is an optimum solution of Problem (1.5), then (\bar{x}, \bar{y}) is an optimum solution of Problem (1.1), with optimum value \bar{x}_0 for the objective function.

Based on this theorem, he has designed a multi-step procedure for solving Problem (1.1).

The procedure starts from a subset Q of C and solves a programming problem

$$(1.6) \quad \text{Max}\{x_0 | (x_0, y) \in G(Q)\},$$

where $G(Q)$ is a set defined by

$$(1.7) \quad G(Q) = \bigcap_{(u_0, u) \in Q} \{(x_0, y) | u_0 x_0 + u'F(y) - u_0 f(y) \leq u'b, y \in S\}.$$

Let an optimum solution of (1.6) be (\bar{x}_0, \bar{y}) . Then, solve the problem

$$(1.8) \quad \text{Min}\{(b - F(\bar{y}))'u | A'u \geq c, u \geq 0\}.$$

Let an optimum solution of (1.8) be \bar{u} . Then, we have:

Theorem 1.2

If (\bar{x}_0, \bar{y}) is an optimum solution of Problem (1.6), it is also an optimum solution of Problem (1.4) if and only if

$$(b - F(\bar{y}))' \bar{u} = \bar{x}_0 - f(\bar{y}).$$

And, if equality holds, we get an optimum solution (\bar{x}, \bar{y}) of Problem (1.1), where \bar{x} is an optimum solution of the linear programming problem (1.5).

On the other hand, if

$$(b - F(\bar{y}))' \bar{u} < \bar{x}_0 - f(\bar{y}),$$

then extend the set Q by adding a certain vertex of the feasible region of Problem (1.8) and/or a certain extremal ray of the convex cone C . And return to Problem (1.6). Repeat the above procedures until an optimum solution of Problem (1.1) is found or Problem (1.6) and hence (1.1) are decided to have no feasible solution or Problem (1.1) is decided to have no finite optimum solution, since Problem (1.8) has no feasible solution.

1.2 Parametric Programming and Primal-Dual Algorithm

KELLEY (1959) has presented a primal-dual algorithm for solving such a type of parametric programming problems as

$$(1.9) \quad \text{Max}\{y'b | y'A \leq c' + \lambda d'\},$$

where A is an (m, n) matrix, b and y are m -vectors, c and d are n -vectors and λ is a scalar parameter.

We assume that an optimum solution \bar{y} of Problem (1.9) for $\lambda = \lambda_0$ is known. And we are going to find an optimum solution of Problem (1.9) for λ less than λ_0 .

First, define an index set S by

$$(1.10) \quad S = \{j | w_j \equiv (c + \lambda_0 d)_j - y'A_j = 0, 1 \leq j \leq n\},$$

where $(c + \lambda_0 d)_j$ is the j -th element of vector $(c + \lambda_0 d)$ and A_j is the j -th column of matrix A .

And then, define the *restricted problem* as follows:

$$(1.11) \quad \text{Min}\{\sigma'b | \sigma'A_j \geq d_j (j \in S)\},$$

where σ is an m -vector.

Now, KELLEY's algorithm is as follows:

Step 1. Solving the restricted problem (1.11)

Solve Problem (1.11). If it has no optimum solution, then Problem (1.9) has no feasible solution for λ less than λ_0 . (*The end.*) Otherwise, let an optimum solution of (1.11) be $\bar{\sigma}$.

Step 2. Finding the bound of θ

Let $\beta_j = d_j - \bar{\sigma}'A_j (1 \leq j \leq n)$.

Find a positive number θ_0 by

$$(1.12) \quad \theta_0 = \begin{cases} \min\{w_j | \beta_j | \beta_j > 0\} \\ \infty (\text{if } \beta_j \leq 0, 1 \leq j \leq n). \end{cases}$$

Step 3. Getting optimum solutions

Step 3.1. An optimum solution of Problem (1.9) for $\lambda = \lambda_0 - \theta$ ($0 \leq \theta \leq \theta_0$), is $\bar{y}' - \theta \bar{\sigma}'$.

Step 3.2. Let \bar{x} be an optimum solution of the dual problem of (1.11). Then, \bar{x} is also an optimum solution of the dual problem of (1.9) for $\lambda = \lambda_0 - \theta$ ($0 \leq \theta \leq \theta_0$).

(This completes one cycle of the algorithm.)

2. A Parametrization of Benders Decomposition

We consider the following parametric programming problem.

[Problem I(λ)] (with variables x and y)

$$\text{Max}\{c'x + f'y \mid Ax + Fy \leq b + \lambda d, 0 \leq x \in R^p, y \in S \subset R^q, \lambda \in R\},$$

where A is an (m, p) matrix, F an (m, q) matrix, b and d m -vectors, c a p -vector, and f a q -vector.

Notice that the variable y appears in linear forms in our problem.

Now, we define several problems and sets corresponding to those in Section 1.1.

[Problem II($y|\lambda$)] (with variable x)

$$\text{Max}\{z_1 = c'x \mid x \geq 0, Ax \leq b + \lambda d - Fy\}.$$

[Problem III ($y|\lambda$)] (with variable u)

$$\text{Min}\{z_2 = u'(b + \lambda d - Fy) \mid A'u \geq c, u \geq 0\}. \quad (\text{The dual problem of Problem II})$$

[Polyhedral Convex Cone C and Set G]

$$C = \{(u_0, u) \mid A'u - cu_0 \geq 0, u \geq 0, u_0 \geq 0\}.$$

$$G = \bigcap_{(u_0, u) \in C} \{(x_0, y) \mid u_0 x_0 + u'Fy - u_0 f'y \leq u'(b + \lambda d), y \in S\}.$$

[Sets Q and $G(Q|\lambda)$]

$$Q = \text{A finite subset of } C.$$

$$G(Q|\lambda) = \bigcap_{(u_0, u) \in Q} \{(x_0, y) \mid u_0 x_0 + u'Fy - u_0 f'y \leq u'(b + \lambda d), y \in S\}.$$

[Problem IV($G(Q|\lambda)$)] (with variables x_0 and y)

$$\text{Max}\{x_0 \mid (x_0, y) \in G(Q|\lambda)\}.$$

Notice that the polyhedral convex cone C has no relation to the parameter λ .

In what follows, we assume that an optimum solution (\bar{x}, \bar{y}) of Problem I(λ) is known for $\lambda = \lambda_0$. (If not, we may use Benders decomposition to find out one.) Also, we can naturally assume, from the above assumption, that we have a set Q and a point $(1, \bar{u}) \in Q$ and we have the relation showing the optimality of \bar{y}, \bar{u} and \bar{x} :

$$\min\{z_2 = u'(b + \lambda_0 d - F\bar{y}) \mid A'u \geq c, u \geq 0\} = \bar{u}'(b + \lambda_0 d - F\bar{y}) = \bar{x}_0 - f'\bar{y} = c'\bar{x},$$

where \bar{x}_0 is the optimum value for the objective function of Problem IV ($G(Q|\lambda_0)$). And we are going to find an optimum solution of Problem I(λ) for λ less than λ_0 . We will deal with two cases corresponding to the kinds of S , namely, the case $S=R^q$ and the case S =a set of discrete points in R^q .

3. The Case $S=R^q$

3.1 Algorithm

In this section, we will show an algorithm for finding an optimum solution of Problem I(λ) in the case $S=R^q$ for λ less than λ_0 , in the knowledge of an optimum solution (\bar{x}, \bar{y}) of Problem I(λ_0).

Step 1. Solving the restricted problem

Based on the optimum solution (\bar{x}_0, \bar{y}) of Problem IV($G(Q|\lambda_0)$), derive its *restricted problem* corresponding to Problem (1.10) of Section 1 and solve it. If it has no optimal solution, then Problem I(λ) has no feasible solution for λ less than λ_0 . (*The end.*) Otherwise, let an optimum solution be $(\bar{\xi}_0, \bar{\eta})$. And let

$$(3.1) \quad x_0(\theta) = \bar{x}_0 - \theta \bar{\xi}_0, \quad y(\theta) = \bar{y} - \theta \bar{\eta}$$

Also, determine the range $[\lambda_0 - \theta_1, \lambda_0]$ of λ where $(x_0(\theta), y(\theta))$ remains optimal for Problem IV($G(Q|\lambda)$), using formula (1.12).

Step 2. Solving Problem I($\lambda_0 - \theta$)

Step 2.1. Derive the *restricted problem* of Problem II($y(\theta)|\lambda_0 - \theta$), in the knowledge of its optimum solution \bar{x} for $\theta=0$ and solve it. Let an optimum solution be $\bar{\xi}$ and let

$$(3.2) \quad x(\theta) = \bar{x} - \theta \bar{\xi}.$$

Then, determine the range $[\lambda_0 - \theta_2, \lambda_0]$ of λ where $x(\theta)$ remains optimal for Problem II($y(\theta)|\lambda$). In this range, \bar{u} remains optimal for Problem III($y(\theta)|\lambda$).

Step 2.2. Let

$$(3.3) \quad \theta_0 = \min \{\theta_1, \theta_2\}.$$

Step 2.3. An optimum solution of Problem I($\lambda_0 - \theta$) for $0 \leq \theta \leq \theta_0$, is $(x(\theta), y(\theta))$.

Step 3. Finding next starting solution

When θ goes out of the range, then $(x(\theta), y(\theta))$ is no longer optimal. Apply Benders decomposition, starting from the present Q and using $y(\theta)$ or \bar{u} if necessary, to find an optimum solution of Problem I($\lambda_0 - \theta_0 - \varepsilon$) where ε is a sufficiently small positive number.

(*This completes one cycle of the algorithm.*)

3.2 Validity of the Algorithm

Proposition 3.1.

If the restricted problem in Step 1 has no optimum solution, then Problem I(λ)

has no feasible solution for λ less than λ_0 .

Proof: By Theorem 1.1, Problem $I(\lambda)$ is not feasible if and only if Problem $IV(G|\lambda)$ is not feasible. Since $Q \subset C$, we have $G(Q) \supset G$. Hence, if $G(Q)$ is empty, then also G .
Q.E.D.

This proposition works in Step 1 as the termination criterion.

Proposition 3.2.

$(x(\theta), y(\theta))$, obtained in Step 1 and Step 2, is an optimum solution of Problem $I(\lambda_0 - \theta)$ for $0 \leq \theta \leq \theta_0$.

Proof: (i) First, we will demonstrate the equality

$$(3.4) \quad x(\theta) + \bar{u}' F y(\theta) - f' y(\theta) = \bar{u}' (b + (\lambda_0 - \theta)d). \quad (0 \leq \theta \leq \theta_0)$$

Let Q_1 be a subset of Q such that $(u_0, u) \in Q_1$ satisfies

$$u_0 \bar{x}_0 + u' F \bar{y} - u_0 f' \bar{y} = u' (b + \lambda_0 d).$$

Of course, $(1, \bar{u}) \in Q_1$ by assumption.

Then, the restricted problem in Step 1 is

$$\text{Min}\{\bar{\xi}_0 | u_0 \bar{\xi}_0 + u' F \bar{\gamma} - u_0 f' \bar{\gamma} \geq u' d, (u_0, u) \in Q_1\}.$$

Its optimum solution $(\bar{\xi}_0, \bar{\gamma})$ must satisfy, for some $(1, u^*) \in Q_1$, the equality

$$\bar{\xi}_0 + u^{*'} F \bar{\gamma} - f' \bar{\gamma} = u^{*'} d.$$

We can assume that this $(1, u^*)$ is $(1, \bar{u})$, without losing generality, because u^* is also an optimum solution of Problem $III(\bar{y}|\lambda_0)$.

Thus, we have

$$\bar{\xi}_0 + \bar{u}' F \bar{\gamma} - f' \bar{\gamma} = \bar{u}' d.$$

Therefore,

$$(\bar{x}_0 - \theta \bar{\xi}_0) + \bar{u}' F(\bar{y} - \theta \bar{\gamma}) - f'(\bar{y} - \theta \bar{\gamma}) = \bar{u}' (b + (\lambda_0 - \theta)d).$$

This shows the equality (3.4).

(ii) Since $y(\theta)$ is an optimum solution of Problem $IV(G(Q)|\lambda_0 - \theta)$ and \bar{u} is an optimum solution of Problem $III(y(\theta)|\lambda_0 - \theta)$ in the range $0 \leq \theta \leq \theta_0$, and we have the equality (3.4), we can conclude that $(x(\theta), y(\theta))$ is optimal for Problem $I(\lambda_0 - \theta)$ ($0 \leq \theta \leq \theta_0$), by Theorem 1.2, $x(\theta)$ being an optimum solution of Problem $II(y(\theta)|\lambda_0 - \theta)$.
Q.E.D.

The above propositions demonstrate the validity of the algorithm.

4. The Discrete Variable Case

4.1 Algorithm

We will show an algorithm for solving the parametric Problem $I(\lambda)$ when S

is a finite set of discrete points in R^q . Similarly to the preceding section, we assume that we have already found an optimum solution (\bar{x}, \bar{y}) of Problem I(λ_0), the set Q , an optimum solution \bar{u} of Problem III($\bar{y}|\lambda_0$) and an optimum solution (\bar{x}_0, \bar{y}) of Problem IV ($G(Q|\lambda_0)$); and we have the equality $\bar{u}'(b + \lambda_0 d - F\bar{y}) = \bar{x}_0 - f'\bar{y}$.

Step 1. Finding the bound of decrease of λ

Step 1.1. Try the parametric analysis of Problem IV($G(Q|\lambda)$) with respect to λ and determine the range $[\lambda_0 - \theta_1, \lambda_0]$ ($\theta_1 \geq 0$) where the solution \bar{y} remains optimal. If there is no feasible solution of Problem IV($G(Q|\lambda)$) for $\lambda < \lambda_0$, then Problem I(λ) has no feasible solution for λ less than λ_0 . (*The end.*)

Step 1.2. Otherwise, by applying the parametric analysis of Problem III($\bar{y}|\lambda$) with respect to λ , determine the range $[\lambda_0 - \theta_2, \lambda_0]$ ($\theta_2 \geq 0$) where the solution \bar{u} remains optimal.

Step 1.3. Let $\theta_0 = \min \{\theta_1, \theta_2\}$.

Step 2. Determining an optimum solution of Problem I(λ)

If $\theta_0 = 0$, go to Step 3. Otherwise, solve Problem II($\bar{y}|\lambda_0 - \theta$) ($0 \leq \theta \leq \theta_0$). Let an optimum solution be $\bar{x} - \theta \bar{\xi}$, where $\bar{\xi}$ is an optimum solution of the restricted problem II($\bar{y}|\lambda_0$). Then, we have an optimum solution $(\bar{x} - \theta \bar{\xi}, \bar{y})$ of Problem I($\lambda_0 - \theta$) for $0 \leq \theta \leq \theta_0$.

Step 3. Finding next starting solution

Apply Benders decomposition, starting from the present Q and using \bar{y} or \bar{u} if necessary, to find an optimum solution of Problem I($\lambda_0 - \theta_0 - \varepsilon$) where ε is a sufficiently small positive number.

(*This completes one cycle of the algorithm.*)

4.2 Validity of the Algorithm

Proposition 4.1.

If Problem IV($G(Q|\lambda)$) has no feasible solution for $\lambda < \lambda_0$ in Step 1.1, then Problem I(λ) has no feasible solution for $\lambda < \lambda_0$.

Proof: The same as Proposition 3.1.

Q.E.D.

Proposition 4.2.

Let the maximum value for the objective function x_0 of Problem IV($G(Q|\lambda_0 - \theta)$) ($0 \leq \theta \leq \theta_0$) be $x_0(\theta)$. Then, $x_0(\theta)$ satisfies

$$x_0(\theta) = \bar{x}_0 - \theta \bar{u}' d.$$

Proof: By the definition of θ_0 , \bar{y} is an optimum solution of Problem IV($G(Q|\lambda_0 - \theta)$) ($0 \leq \theta \leq \theta_0$). Therefore, for every $(u_0, u) \in Q$, we have

$$u_0 x_0(\theta) + u' F \bar{y} - u_0 f' \bar{y} \leq u'(b + (\lambda_0 - \theta)d).$$

And, since $(1, \bar{u}) \in Q$, we have

$$x_0(\theta) + \bar{u}' F \bar{y} - f' \bar{y} \leq \bar{u}'(b + (\lambda_0 - \theta)d).$$

But, for $\theta = 0$, from the optimality of (\bar{x}_0, \bar{y}) and $(1, \bar{u})$, we have

$$\bar{x}_0 + \bar{u}'F\bar{y} - f'\bar{y} = \bar{u}'(b + \lambda_0 d).$$

Hence, the following relation holds:

$$x_0(\theta) \leq \bar{x}_0 - \theta \bar{u}'d.$$

Now, we demonstrate the equality. Suppose the contrary. Then, for some $\theta (0 \leq \theta \leq \theta_0)$, there exists at least a $(1, \bar{u}) \in Q$, such that

$$(4.1) \quad x_0(\theta) = -\bar{u}'F\bar{y} + f'\bar{y} - \bar{u}'(b + (\lambda_0 - \theta)d) > -\bar{u}'F\bar{y} + f'\bar{y} - \bar{u}'(b + (\lambda_0 - \theta)d).$$

But, since \bar{u} is a feasible solution of Problem III($\bar{y}|\lambda_0 - \theta$) and \bar{u} is an optimum solution, we have

$$\bar{u}'(b + (\lambda_0 - \theta)d - F\bar{y}) \leq \bar{u}'(b + (\lambda_0 - \theta)d - F\bar{y}).$$

This contradicts (4.1)

Q.E.D.

Proposition 4.3.

The solution $(\bar{x} - \theta \bar{\xi}, \bar{y})$ in Step 2 is an optimum solution of Problem I(λ) for $\lambda = \lambda_0 - \theta (0 \leq \theta \leq \theta_0)$.

Proof: Proposition 4.2 means

$$x_0(\theta) - f'\bar{y} = \bar{u}'(b + (\lambda_0 - \theta)d - F\bar{y}).$$

By Theorem 1.2, this shows the optimality of $(\bar{x} - \theta \bar{\xi}, \bar{y})$.

Q.E.D.

The above propositions demonstrate the validity of the algorithm.

Concluding Remarks

Benders decomposition is recognized as an excellent partitioning procedure for solving such a mixed type problem that involves both structured variables and non-structured variables. Our algorithm deals with a parametrization of Benders decomposition in the descending value of the parameter. But we can easily modify the algorithm to be valid for the ascending case. TONE (forthcoming) shows an example of applications of algorithms in this paper.

REFERENCES

- J. F. BENDERS (1962): Partitioning procedures for solving mixed-variables programming problems, *Numerische Mathematik*, **4**, pp. 238-252.
- J. E. KELLEY, JR. (1959): Parametric programming and the primal-dual algorithm, *Operations Research*, **7**, pp. 327-334.
- K. TONE (forthcoming): On optimal pattern flows, submitted to *Journal of the Operations Research Society of Japan*.