| Title            | Generalized harmonic analysis of functions of two variables  |
|------------------|--|
| Sub Title        |  |
| Author           | 松岡, 勝男(Matsuoka, Katsuo)   |
| Publisher        | 慶応義塾大学工学部  |
| Publication year | 1980   |
| Jtitle           | Keio engineering reports Vol.33, No.6 (1980. 5) ,p.67- 81  |
| JaLC DOI         |  |
| Abstract         | On the basis of the Wiener formula of functions of two variables under a restricted limit process, a new approach to the generalized harmonic analysis of functions of two variables is shown. |
| Notes            |  |
| Genre            | Departmental Bulletin Paper  |
| URL              | https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=KO50001004-00330006-<br>0067  |

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

# GENERALIZED HARMONIC ANALYSIS OF FUNCTIONS OF TWO VARIABLES

ΚΑΤSUO ΜΑΤSUOKA

Dept. of Mathematics, Keio University, Yokohama 223, Japan

(Received January, 24, 1980)

### ABSTRACT

On the basis of the Wiener formula of functions of two variables under a restricted limit process, a new approach to the generalized harmonic analysis of functions of two variables is shown.

### §1. Introduction

Wiener [10] established the generalized harmonic analysis for the analysis of paths of the Brownian motion. Wiener and A. C. Berry also proved the case of functions of two variables (*Wiener* [10]). As is well-known, it is based on the so-called Wiener formula. They use the circular mean concerning the double limit process. It seems to be something restricted.

Now, it is worthwhile to consider a possibility of approaching this problem by a more relaxed limit process. Recently, Anzai, Koizumi and Matsuoka [1] proved the two-dimensional Wiener formula under a restricted rectangular mean concerning the double limit process. In this paper, we shall establish the generalized harmonic analysis of functions of two variables under this restricted limit process.

The proofs can be done along the similar lines as in Wiener [11].

## §2. Definitions and theorems

First, we introduce the following class of functions.

DEFINITION 1. By  $W(R^2)$ , we denote the class of functions such that  $f(x_1, x_2) \in L^2_{loc}(R^2)$  and

$$\frac{1}{4ST} \int_{-r}^{r} \int_{-s}^{s} |f(s,t)|^2 ds dt$$
 (2.1)

is bounded in S, T > 0.

Then we shall prove

THEOREM 1. If  $f(x_1, x_2)$  belongs to the class  $W(R^2)$ , then

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{|f(s,t)|^2}{(1+s^2)(1+t^2)} \, ds dt < \infty \,. \tag{2.2}$$

Next, we introduce the double generalized Fourier transform due to Wiener [10, 11]. This is defined by

$$s(u, v; f) = \lim_{A \to \infty} \frac{1}{2\pi} \left[ \int_{1}^{A} + \int_{-A}^{-1} \right] \left[ \int_{1}^{A} + \int_{-A}^{-1} \right] f(s, t) \frac{e^{-ius}}{-is} \frac{e^{-ivt}}{-it} \, ds dt \\ + \lim_{A \to \infty} \frac{1}{2\pi} \left[ \int_{1}^{A} + \int_{-A}^{-1} \right] \int_{-1}^{1} f(s, t) \frac{e^{-ius} - 1}{-is} \frac{e^{-ivt}}{-it} \, ds dt \\ + \lim_{A \to \infty} \frac{1}{2\pi} \int_{-1}^{1} \left[ \int_{1}^{A} + \int_{-A}^{-1} \right] f(s, t) \frac{e^{-ius}}{-is} \frac{e^{-ivt} - 1}{-it} \, ds dt \\ + \frac{1}{2\pi} \int_{-1}^{1} \int_{-1}^{1} f(s, t) \frac{e^{-ius} - 1}{-is} \frac{e^{-ivt} - 1}{-it} \, ds dt .$$
(2.3)

If  $f(x_1, x_2)$  belongs to the class  $W(\mathbb{R}^2)$ , then by Theorem 1 and the Plancherel theorem, the double generalized Fourier transform s(u, v; f) is defined and we have

$$\Delta_{\iota,\eta} s(u,v;f) = s(u+\varepsilon,v+\eta;f) - s(u-\varepsilon,v+\eta;f) - s(u+\varepsilon,v-\eta;f) + s(u-\varepsilon,v-\eta;f)$$
$$= \lim_{A \to \infty} \frac{1}{2\pi} \int_{-A}^{A} \int_{-A}^{A} f(s,t) \frac{2\sin\varepsilon s}{s} \frac{2\sin\eta t}{t} e^{-i(us-vt)} ds dt$$
(2.4)

and

$$\frac{1}{16\pi^2 \varepsilon \eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |\Delta_{\epsilon, \eta} s(u, v; f)|^2 du dv$$
  
=  $\frac{1}{\pi^2 \varepsilon \eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |f(s, t)|^2 \frac{\sin^2 \varepsilon s}{s^2} \frac{\sin^2 \eta t}{t^2} ds dt$ . (2.5)

On the other hand, Anzai, Koizumi and Matsuoka [1] proved the following twodimensional Wiener formula in the first quadrant  $R_{+}^2$  of the plane.

THEOREM 2. Suppose that  $f(x_1, x_2) \ge 0$  in  $(x_1, x_2) \in R_+^2$ ,  $f(x_1, x_2) \in L^1_{loc}(R_+^2)$  and

$$\frac{1}{ST} \int_0^T \int_0^s f(s, t) ds dt$$
(2.6)

is bounded in S, T>0. Then the limit relations

$$\lim_{T \to \infty} \frac{1}{C_1 C_2 T^2} \int_0^{C_2 T} \int_0^{C_1 T} f(x_1, x_2) dx_1 dx_2 = A \quad (\forall C_1, C_2 > 0)$$
(2.7)

and

$$\lim_{\varepsilon \to 0} \frac{4C_1C_2}{\pi^2 \varepsilon^2} \int_0^\infty \int_0^\infty f(x_1, x_2) \frac{\sin^2 \varepsilon C_1^{-1} x_1}{x_1^2} \frac{\sin^2 \varepsilon C_2^{-1} x_2}{x_2^2} dx_1 dx_2 = A \quad (\forall C_1, C_2 > 0)$$
(2.8)

are equivalent in the sense that if either of the limits (2.7) or (2.8) exists, then the other limit exists and assumes the same value.

By the way, if we put S, T,  $\varepsilon$ ,  $\eta$  and C instead of  $C_1T$ ,  $C_2T$ ,  $\varepsilon C_1^{-1}$ ,  $\varepsilon C_2^{-1}$  and  $C_1/C_2$  respectively, then S=CT and  $\eta=C\varepsilon$  hold. Hence, (2.7) and (2.8) of Theorem 2 are equivalent to the statements

$$\lim_{s, \tau \to \infty} \frac{1}{ST} \int_0^\tau \int_0^s f(s, t) ds dt$$

exists and has the same limit for every positive constant C whenever S and T tend to infinity in such a way that S=CT " and

$$\lim_{\iota,\eta\to 0}\frac{4}{\pi^2\varepsilon\eta}\int_0^\infty\int_0^\infty f(s,t)\frac{\sin^2\varepsilon s}{s^2}\frac{\sin^2\eta t}{t^2}\,dsdt$$

exists and has the same limit for every positive constant C whenever  $\varepsilon$  and  $\eta$  tend to zero in such a way that  $\eta = C\varepsilon$  " respectively.

For the sake of simplicity, we shall from now on use the notations  $\mathcal{R}_1$  and  $\mathcal{R}_2$  instead of denoting the above limit processes respectively. And we shall refer to these limits as the restricted limits. Therefore, using the notations  $\mathcal{R}_1$  and  $\mathcal{R}_2$ , Theorem 2 is rewritten as follows:

THEOREM 2'. If  $f(x_1, x_2)$  satisfies the hypotheses of Theorem 2, and either of the limits

$$\mathcal{R}_{1} - \lim_{S,T \to \infty} \frac{1}{ST} \int_{0}^{T} \int_{0}^{S} f(s,t) ds dt$$
(2.9)

or

$$\mathcal{R}_{2} - \lim_{\epsilon, \eta \to 0} \frac{4}{\pi^{2} \epsilon \eta} \int_{0}^{\infty} \int_{0}^{\infty} f(s, t) \frac{\sin^{2} \epsilon s}{s^{2}} \frac{\sin^{2} \eta t}{t^{2}} \, ds dt$$
(2.10)

exists, then the other limit exists and assumes the same value.

From this theorem and (2.5), we get

THEOREM 3. If  $f(x_1, x_2)$  belongs to the class  $W(R^2)$ , then we have

$$\mathcal{R}_{1} \cdot \lim_{S,T \to \infty} \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} |f(s,t)|^{2} ds dt$$
$$= \mathcal{R}_{2} \cdot \lim_{\epsilon,\eta \to 0} \frac{1}{16\pi^{2}\epsilon\eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |\Delta_{\epsilon,\eta} s(u,v;f)|^{2} du dv, \qquad (2.11)$$

in the sense that if either side of (2.11) exists, the other side exists and assumes the same value.

Now, we introduce the following two classes of functions which are due to Wiener [10, 11].

DEFINITION 2. By  $S(\mathbb{R}^2)$ , we denote the class of functions such that  $f(x_1, x_2) \in W(\mathbb{R}^2)$  and

$$\phi(x_1, x_2; f) = \mathcal{R}_1 - \lim_{s. \tau \to \infty} \frac{1}{4ST} \int_{-\tau}^{\tau} \int_{-s}^{s} f(x_1 + s, x_2 + t) \overline{f(s, t)} ds dt$$
(2.12)

exists for all  $(x_1, x_2) \in \mathbb{R}^2$ .

DEFINITION 3. By  $S'(R^2)$ , we denote the class of functions such that  $f(x_1, x_2) \in S(R^2)$  and  $\phi(x_1, x_2; f)$  is continuous on  $R^2$ .

 $S'(R^2)$  is a proper subclass of  $S(R^2)$ . In other words, there is a function  $\phi(x_1, x_2; f)$  defined by (2.12) which is not continuous on  $R^2$ . For example, take  $f(x_1, x_2) = \exp \{i(x_1^2 + x_2^2)\}$ . Then

$$\begin{aligned} \phi(x_1, x_2; f) &= \mathcal{R}_1 \cdot \lim_{s, \tau \to \infty} \frac{1}{4ST} \int_{-\tau}^{\tau} \int_{-s}^{s} e^{i((x_1+s)^2 + (x_2+t)^2)} e^{-i(s^2+t^2)} ds dt \\ &= \begin{cases} 1 & ((x_1, x_2) = (0, 0)) \\ 0 & (\text{elsewhere}) \end{cases}. \end{aligned}$$

Here, we shall consider the properties of functions of  $S(R^2)$  and  $S'(R^2)$ .

THEOREM 4. If  $f(x_1, x_2)$  belongs to the class  $S(R^2)$ , then

$$|\phi(x_1, x_2; f)| \le \phi(0, 0; f) \tag{2.13}$$

for all  $(x_1, x_2) \in \mathbb{R}^2$ .

THEOREM 5. Suppose that  $f(x_1, x_2)$  belongs to the class  $S(R^2)$  and  $\phi(x_1, x_2; f)$  is continuous at  $(x_1, x_2) = (0, 0)$ . Then  $\phi(x_1, x_2; f)$  is continuous at every point of  $R^2$  and  $f(x_1, x_2)$  belongs to the class  $S'(R^2)$ .

THEOREM 6. If  $f(x_1, x_2)$  belongs to the class  $S(\mathbb{R}^2)$ , then

$$\phi(x_1, x_2; f) = \mathcal{R}_2 \lim_{\epsilon, \eta \to 0} \frac{1}{16\pi^2 \epsilon \eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{i(ux_1 + vx_2)} |\Delta_{\epsilon, \eta} s(u, v; f)|^2 du dv .$$
(2.14)

THEOREM 7. If  $f(x_1, x_2)$  belongs to the class  $S(R^2)$ , then it will belong to the class  $S'(R^2)$  when and only when both

$$\lim_{A \to \infty} \mathcal{R}_2 \overline{\lim_{\epsilon, \eta \to 0}} \frac{1}{16\pi^2 \epsilon \eta} \int_{-\infty}^{\infty} \left[ \int_{A}^{\infty} + \int_{-\infty}^{-A} \right] |\Delta_{\epsilon, \eta} s(u, v; f)|^2 du dv = 0$$
(2.15)

and

$$\lim_{A \to \infty} \mathcal{R}_2 \overline{\lim_{\epsilon, \eta \to 0}} \frac{1}{16\pi^2 \varepsilon \eta} \left[ \int_A^{\infty} + \int_{-\infty}^{-A} \right] \int_{-\infty}^{\infty} |\Delta_{\epsilon, \eta} S(u, v; f)|^2 du dv = 0$$
(2.16)

are true.

Finally, we shall show Theorem 8 below concerning the spectral representation of the correlation function  $\phi(x_1, x_2; f)$  of  $f(x_1, x_2)$ . For this, we first observe the following two well-known theorems for positive definite functions. Now, let  $\mathcal{B}_2$  be the Borel field in  $\mathbb{R}^2$ .

Theorem (Bochner's representation theorem).\* If  $\phi(x_1, x_2)$  is a positive definite function, then there exists a measure  $\mu$  on  $(R^2, \mathcal{B}_2)$  such that  $\mu(R^2) = \phi(0, 0)$ , and

$$\phi(x_1, x_2) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{i(\xi_1 x_1 + \xi_2 x_2)} d\mu(\xi_1, \xi_2)$$
(2.17)

holds.

Theorem (Lévy's inversion formula). The measure  $\mu$  of (2.17) of Bochner's representation theorem is represented, by  $\phi(x_1, x_2)$ , as follows: If  $I = (\alpha_1, \alpha_2] \times (\beta_1, \beta_2]$  is a finite continuity interval of  $\mu$ , then we have

$$\mu(I) = \lim_{A \to \infty} \frac{1}{2\pi} \int_{-A}^{A} \int_{-A}^{A} \frac{e^{-i\alpha_2 x_1} - e^{-i\alpha_1 x_1}}{-ix_1} \frac{e^{-i\beta_2 x_2} - e^{-i\beta_1 x_2}}{-ix_2} \phi(x_1, x_2) dx_1 dx_2 .$$
(2.18)

Thus, the positive definite function  $\phi(x_1, x_2)$  determines uniquely the measure  $\mu$ .

We second observe the following definition.

DEFINITION 4. A function  $\Lambda(u, v)$  is called monotone increasing if

$$\Lambda(u+h,v+k) - \Lambda(u,v+k) - \Lambda(u+h,v) + \Lambda(u,v) \ge 0$$
(2.19)

for all  $h, k \ge 0$ .

Then we can state

THEOREM 8. If  $f(x_1, x_2)$  belongs to the class  $S'(R^2)$ , then  $\phi(x_1, x_2; f)$  is positive definite. Thus, there exists a monotone increasing function  $\Lambda(u, v)$  such that  $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |d\Lambda(u, v)| = \phi(0, 0; f)$ , and

$$\phi(x_1, x_2; f) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{i(ux_1 + vx_2)} dA(u, v)$$
(2.20)

holds. Moreover, if  $(\alpha_1, \alpha_2] \times (\beta_1, \beta_2]$  is a finite continuity interval of the interval function generated by  $\Lambda(u, v)$ , then we have

$$\begin{aligned}
\Lambda(\alpha_2, \beta_2) - \Lambda(\alpha_1, \beta_2) - \Lambda(\alpha_2, \beta_1) + \Lambda(\alpha_1, \beta_1) \\
= \lim_{A \to \infty} \frac{1}{2\pi} \int_{-A}^{A} \int_{-A}^{A} \frac{e^{-i\alpha_2 x_1} - e^{-i\alpha_1 x_1}}{-ix_1} \frac{e^{-i\beta_2 x_2} - e^{-i\beta_1 x_2}}{-ix_2} \phi(x_1, x_2; f) dx_1 dx_2. \quad (2.21)
\end{aligned}$$

<sup>\*</sup> Here, by positive definiteness, we mean the function under consideration to be continuous.

# §3. Proofs of theorems

First, we prove Theorem 1.

Proof of THEOREM 1. If we integrate by parts for each variable, then

$$\begin{split} \int_{-B}^{B} \int_{-A}^{A} \frac{|f(s,t)|^{2}}{(1+s^{2})(1+t^{2})} ds dt \\ &= \frac{1}{(1+A^{2})(1+B^{2})} \int_{-B}^{B} \int_{-A}^{A} |f(\xi_{1},\xi_{2})|^{2} d\xi_{1} d\xi_{2} \\ &+ \frac{1}{1+A^{2}} \int_{0}^{B} \frac{4t^{2}}{(1+t^{2})^{2}} dt \frac{1}{2t} \int_{-t}^{t} \int_{-A}^{A} |f(\xi_{1},\xi_{2})|^{2} d\xi_{1} d\xi_{2} \\ &+ \frac{1}{1+B^{2}} \int_{0}^{A} \frac{4s^{2}}{(1+s^{2})^{2}} ds \frac{1}{2s} \int_{-B}^{B} \int_{-s}^{s} |f(\xi_{1},\xi_{2})|^{2} d\xi_{1} d\xi_{2} \\ &+ \int_{0}^{B} \int_{0}^{A} \frac{4s^{2}}{(1+s^{2})^{2}} \frac{4t^{2}}{(1+t^{2})^{2}} ds dt \frac{1}{4st} \int_{-t}^{t} \int_{-s}^{s} |f(\xi_{1},\xi_{2})|^{2} d\xi_{1} d\xi_{2} \\ &\leq \left\{ \frac{4AB}{(1+A^{2})(1+B^{2})} + \frac{8A}{1+A^{2}} \tan^{-1}B + \frac{8B}{1+B^{2}} \tan^{-1}A + 16 \tan^{-1}A \tan^{-1}B \right\} \\ &\cdot \sup_{0 \leq S, T < \infty} \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} |f(s,t)|^{2} ds dt \\ &\leq \left\{ 1 + 4\pi + 4\pi^{2} \right\} \cdot \sup_{0 < S, T < \infty} \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} |f(s,t)|^{2} ds dt < \infty \,. \end{split}$$

Thus, (2.2) is established.

Next, in order to prove Theorems 4, 5 and 6, we show several lemmas which are due to Wiener [11].

LEMMA 1. If  $\phi(x_1, x_2; f)$  defined by (2.12) exists at  $(x_1, x_2) = (0, 0)$ , then for any real numbers a, b,

$$\mathcal{R}_{1} \lim_{s, \tau \to \infty} \frac{1}{4ST} \int_{-\tau}^{T} \int_{-s}^{s} |f(s+a, t+b)|^{2} ds dt$$
  
=  $\mathcal{R}_{1} \lim_{s, \tau \to \infty} \frac{1}{4ST} \int_{-\tau}^{T} \int_{-s}^{s} |f(s, t)|^{2} ds dt$ . (3.1)

*Proof.* Let us notice that if S > |a|, T > |b|, then

$$\begin{split} & \left(1 - \frac{|a|}{S}\right) \left(1 - \frac{|b|}{T}\right) \frac{1}{4(S - |a|)(T - |b|)} \int_{-T + |b|}^{T - |b|} \int_{-S + |a|}^{S - |a|} |f(s, t)|^2 ds dt \\ & \leq \frac{1}{4ST} \int_{-T + b}^{T + b} \int_{-S + a}^{S + a} |f(s, t)|^2 ds dt = \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} |f(s + a, t + b)|^2 ds dt \\ & \leq \left(1 + \frac{|a|}{S}\right) \left(1 + \frac{|b|}{T}\right) \frac{1}{4(S + |a|)(T + |b|)} \int_{-T - |b|}^{T + |b|} \int_{-S - |a|}^{S + |a|} |f(s, t)|^2 ds dt \,. \end{split}$$

Thus, since both the first and the last of these expressions tend to  $\phi(0,0; f)$  as

S,  $T \rightarrow \infty(\mathcal{R}_1)$ , we have (3.1).

Now, we have

$$f(x_1+s, x_2+t)\overline{f(s,t)} = \frac{1}{4} \{ |f(x_1+s, x_2+t) + f(s,t)|^2 - |f(x_1+s, x_2+t) - f(s,t)|^2 + i|f(x_1+s, x_2+t) + if(s,t)|^2 - i|f(x_1+s, x_2+t) - if(s,t)|^2 \} .$$
(3.2)

Using this, we can show

LEMMA 2. If  $f(x_1, x_2)$  belongs to the class  $S(\mathbb{R}^2)$ , then for any real numbers a, b,

$$\mathcal{R}_{1} - \lim_{s, T \to \infty} \frac{1}{4ST} \int_{-T}^{T} \int_{-s}^{s} f(x_{1} + s + a, x_{2} + t + b) \overline{f(s + a, t + b)} ds dt$$
  
=  $\mathcal{R}_{1} - \lim_{s, T \to \infty} \frac{1}{4ST} \int_{-T}^{T} \int_{-s}^{s} f(x_{1} + s, x_{2} + t) \overline{f(s, t)} ds dt$ . (3.3)

**Proof.** Since  $f(x_1, x_2)$  belongs to the class  $S(\mathbb{R}^2)$ , by (3.2) and Lemma 1,

$$\begin{split} \mathcal{R}_{1}\text{-}\lim_{S,T\to\infty} \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} f(x_{1}+s+a, x_{2}+t+b) \overline{f(s+a,t+b)} ds dt \\ &= \frac{1}{4} \mathcal{R}_{1}\text{-}\lim_{S,T\to\infty} \frac{1}{4ST} \int_{-T+b}^{T+b} \int_{-S+a}^{S+a} \{|f(x_{1}+s, x_{2}+t)+f(s,t)|^{2} \\ &- |f(x_{1}+s, x_{2}+t)-f(s,t)|^{2} + i|f(x_{1}+s, x_{2}+t)+if(s,t)|^{2} \\ &- i|f(x_{1}+s, x_{2}+t)-f(s,t)|^{2} \} ds dt \\ &= \mathcal{R}_{1}\text{-}\lim_{S,T\to\infty} \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} f(x_{1}+s, x_{2}+t) \overline{f(s,t)} ds dt \; . \end{split}$$

Thus, the lemma is established.

Furthermore, we have for any real or complex number w such as |w|=1,

$$|f(x_1+s, x_2+t) + wf(s, t)|^2 = |f(x_1+s, x_2+t)|^2 + |f(s, t)|^2 + \overline{w}f(x_1+s, x_2+t)\overline{f(s, t)} + w\overline{f(x_1+s, x_2+t)}f(s, t), \quad (3.4)$$

from which we have

LEMMA 3. If  $f(x_1, x_2)$  belongs to the class  $S(R^2)$ , then for any real or complex number w such as |w|=1,

$$\mathcal{R}_{1} - \lim_{S,T \to \infty} \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} |f(x_{1}+s, x_{2}+t) + wf(s,t)|^{2} ds dt$$
(3.5)

exists for all  $(x_1, x_2) \in \mathbb{R}^2$ .

On the other hand, we can obtain

### Katsuo Matsuoka

LEMMA 4. If  $f(x_1, x_2)$  belongs to the class  $W(R^2)$ , then we have for any  $x_1 \in (-\infty, \infty)$ ,

$$\mathcal{R}_{2} - \lim_{\epsilon,\eta \to 0} \frac{1}{16\pi^{2} \varepsilon \eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |\Delta_{\epsilon,\eta} s_{x_{1}}(u,v;f) - e^{iux_{1}} \Delta_{\epsilon,\eta} s(u,v;f)|^{2} du dv = 0$$
(3.6)

and for any  $x_2 \in (-\infty, \infty)$ ,

$$\mathcal{R}_{2} - \lim_{\epsilon,\eta \to 0} \frac{1}{16\pi^{2}\epsilon\eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |\Delta_{\epsilon,\eta} s_{x_{2}}(u,v;f) - e^{ivx_{2}} \Delta_{\epsilon,\eta} s(u,v;f)|^{2} du dv = 0, \qquad (3.7)$$

where

$$\Delta_{\epsilon,\eta} s_{x_1}(u,v;f) = \lim_{A \to \infty} \frac{1}{2\pi} \int_{-A}^{A} \int_{-A}^{A} f(x_1+s,t) \frac{2\sin\epsilon s}{s} \frac{2\sin\eta t}{t} e^{-i(us+vt)} ds dt \qquad (3.8)$$

and

$$\Delta_{\varepsilon,\eta} s_{x_2}(u,v;f) = \lim_{A \to \infty} \frac{1}{2\pi} \int_{-A}^{A} \int_{-A}^{A} f(s, x_2 + t) \frac{2\sin\varepsilon s}{s} \frac{2\sin\eta t}{t} e^{-i(us+vt)} ds dt , \quad (3.9)$$

respectively.

Before proving the lemma, we observe the following: If  $f(x_1, x_2)$  belongs to the class  $W(R^2)$ , then in a way similar to the proof of Theorem 1,

$$\frac{1}{2T} \int_{-T}^{T} dt \int_{-A}^{A} \frac{|f(x_1, t)|^2}{1 + {x_1}^2} dx_1 \leq (1 + 2\pi) \cdot \sup_{0 < S, T < \infty} \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} |f(s, t)|^2 ds dt < \infty.$$

It follows, therefore, that

$$\overline{\lim_{T \to \infty}} \frac{1}{2T} \int_{-T}^{T} dt \int_{-\infty}^{\infty} \frac{|f(x_1, t)|^2}{1 + {x_1}^2} dx_1 < \infty .$$
(3.10)

Similarly,

$$\overline{\lim_{S \to \infty}} \frac{1}{2S} \int_{-S}^{S} ds \int_{-\infty}^{\infty} \frac{|f(s, x_2)|^2}{1 + x_2^2} dx_2 < \infty .$$
(3.11)

Proof. Let us notice that

$$\left|\frac{2\sin\varepsilon(\xi-x)}{\xi-x} - \frac{2\sin\varepsilon\xi}{\xi}\right| \leq \begin{cases} \frac{16\varepsilon|x|}{|\xi|+|x|} & (|\xi|>2|x|)\\ 4\varepsilon & (|\xi|\leq 2|x|) . \end{cases}$$
(3.12)

As for detailed calculations, refer to Wiener [11].

Now, since

$$\begin{aligned} \Delta_{\epsilon,\gamma} s_{x_1}(u,v;f) &- e^{iux_1} \Delta_{\epsilon,\gamma} s(u,v;f) \\ &= \lim_{A \to \infty} \frac{1}{2\pi} \int_{-A}^{A} \int_{-A}^{A} f(s,t) \bigg[ \frac{2\sin \varepsilon(s-x_1)}{s-x_1} - \frac{2\sin \varepsilon s}{s} \bigg] \frac{2\sin \eta t}{t} e^{-i(u(s-x_1)+vt)} ds dt \,, \end{aligned}$$

it follows from the Plancherel theorem and (3.12) that

$$\begin{split} \overline{\lim_{\epsilon \to 0}} & \frac{1}{16\pi^2 C \varepsilon^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |\Delta_{\epsilon, C\epsilon} s_{x_1}(u, v ; f) - e^{iux_1} \Delta_{\epsilon, C\epsilon} s(u, v ; f)|^2 du dv \\ & \leq \overline{\lim_{\epsilon \to 0}} & \frac{(8|x_1|)^2}{\pi} \varepsilon \cdot \frac{1}{\pi C \varepsilon} \int_{-\infty}^{\infty} \frac{\sin^2 C \varepsilon t}{t^2} dt \int_{|s| > 2|x_1|} \frac{|f(s, t)|^2}{(|s| + |x_1|)^2} ds \\ & + \overline{\lim_{\epsilon \to 0}} & \frac{4}{\pi} \varepsilon \cdot \frac{1}{\pi C \varepsilon} \int_{-\infty}^{\infty} \frac{\sin^2 C \varepsilon t}{t^2} dt \int_{|s| \le 2|x_1|} |f(s, t)|^2 ds \,. \end{split}$$

Consequently, using the one-sided Wiener formula of Koizumi [6], the last expression does not exceed, for every positive constant C,

$$\begin{split} \overline{\lim_{\varepsilon \to 0}} \ O(\varepsilon) \cdot \overline{\lim_{T \to \infty}} \frac{1}{2T} \int_{-T}^{T} dt \int_{|s| > 2! x_1|} \frac{|f(s, t)|^2}{(|s| + |x_1|)^2} ds \\ + \overline{\lim_{\varepsilon \to 0}} \ O(\varepsilon) \cdot \overline{\lim_{T \to \infty}} \frac{1}{2T} \int_{-T}^{T} dt \int_{|s| \le 2! x_1|} |f(s, t)|^2 ds \,, \end{split}$$

which turns out to be 0 by (3.10), and (3.6) is proved. Similarly, (3.7) is proved by (3.11).

Moreover, we can easily obtain

LEMMA 5. If  $f(x_1, x_2)$  belongs to the class  $W(R^2)$ , then we have for any  $(x_1, x_2) \in R^2$ ,

$$\mathcal{R}_{2}-\lim_{\epsilon,\eta\to 0}\frac{1}{16\pi^{2}\varepsilon\eta}\int_{-\infty}^{\infty}\int_{-\infty}^{\infty}|\Delta_{\epsilon,\eta}s_{x_{1},x_{2}}(u,v;f) -e^{i(ux_{1}+vx_{2})}\Delta_{\epsilon,\eta}s(u,v;f)|^{2}dudv=0, \qquad (3.13)$$

where

$$\Delta_{\epsilon,\eta} s_{x_1,x_2}(u,v;f) = \lim_{A \to \infty} \frac{1}{2\pi} \int_{-A}^{A} \int_{-A}^{A} f(x_1+s,x_2+t) \frac{2\sin\varepsilon s}{s} \frac{2\sin\eta t}{t} e^{-i(us+vt)} ds dt.$$
(3.14)

*Proof.* First, we have

$$\begin{split} \Delta_{\epsilon,\eta} s_{x_1, x_2}(u, v; f) &= e^{i(ux_1 + vx_2)} \Delta_{\epsilon,\eta} s(u, v; f) \\ &= \{\Delta_{\epsilon,\eta} s_{x_1, x_2}(u, v; f) - e^{ivx_2} \Delta_{\epsilon,\eta} s_{x_1}(u, v; f) \\ &- e^{iux_1} \Delta_{\epsilon,\eta} s_{x_2}(u, v; f) + e^{i(ux_1 + vx_2)} \Delta_{\epsilon,\eta} s(u, v; f)\} \\ &+ \{e^{ivx_2} \Delta_{\epsilon,\eta} s_{x_1}(u, v; f) - e^{i(ux_1 + vx_2)} \Delta_{\epsilon,\eta} s(u, v; f)\} \\ &+ \{e^{iux_1} \Delta_{\epsilon,\eta} s_{x_2}(u, v; f) - e^{i(ux_1 + vx_2)} \Delta_{\epsilon,\eta} s(u, v; f)\} \\ &= \Delta_1(u, v) + \Delta_2(u, v) + \Delta_8(u, v), \quad \text{say.} \end{split}$$

Here  $\Delta_{\iota,\eta}s_{x_1}(u, v; f)$  and  $\Delta_{\iota,\eta}s_{x_2}(u, v; f)$  are defined respectively by (3.8) and (3.9) of Lemma 4. Now, since

### Katsuo Matsuoka

$$\Delta_{1}(u, v) = \lim_{A \to \infty} \frac{1}{2\pi} \int_{-A}^{A} \int_{-A}^{A} f(s, t) \left[ \frac{2 \sin \varepsilon(s - x_{1})}{s - x_{1}} - \frac{2 \sin \varepsilon s}{s} \right]$$
$$\cdot \left[ \frac{2 \sin \eta (t - x_{2})}{t - x_{2}} - \frac{2 \sin \eta t}{t} \right] e^{-i(u(s - x_{1}) + v(t - x_{2}))} ds dt ,$$

it follows from the Plancherel theorem and (3.12) that

$$\begin{split} & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |\Delta_{1}(u, v)|^{2} du dv \\ & \leq (16\varepsilon |x_{1}|)^{2} (16\eta |x_{2}|)^{2} \int_{|t|>2|x_{2}|} \int_{|s|>2|x_{1}|} \frac{|f(s, t)|^{2}}{(|s|+|x_{1}|)^{2} (|t|+|x_{2}|)^{2}} ds dt \\ & + (16\varepsilon |x_{1}|)^{2} (4\eta)^{2} \int_{|t|\leq 2|x_{2}|} \int_{|s|>2|x_{1}|} \frac{|f(s, t)|^{2}}{(|s|+|x_{1}|)^{2}} ds dt \\ & + (4\varepsilon)^{2} (16\eta |x_{2}|)^{2} \int_{|t|>2|x_{2}|} \int_{|s|\leq 2|x_{1}|} \frac{|f(s, t)|^{2}}{(|t|+|x_{2}|)^{2}} ds dt \\ & + (4\varepsilon)^{2} (4\eta)^{2} \int_{|t|\leq 2|x_{2}|} \int_{|s|\leq 2|x_{1}|} |f(s, t)|^{2} ds dt \, . \end{split}$$

Consequently, by Theorem 1,

$$\mathcal{R}_2 - \lim_{\epsilon, \eta \to 0} \frac{1}{16\pi^2 \varepsilon \eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |\Delta_1(u, v)|^2 du dv = 0$$

Also, by Lemma 4,

$$\mathcal{R}_{2} - \lim_{\iota,\eta\to 0} \frac{1}{16\pi^{2}\varepsilon\eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |\Delta_{2}(u,v)|^{2} du dv = 0$$

and

$$\mathcal{R}_2 - \lim_{\varepsilon, \eta \to 0} \frac{1}{16\pi^2 \varepsilon \eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |\Delta_3(u, v)|^2 du dv = 0,$$

respectively. Thus, (3.13) is proved.

Therefore, we have

LEMMA 6. If  $f(x_1, x_2)$  belongs to the class  $S(R^2)$ , then for any real or complex number w such as |w|=1,

$$\mathcal{R}_{1} - \lim_{\boldsymbol{\mathcal{S}},\boldsymbol{T}\to\infty} \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} |f(x_{1}+s, x_{2}+t) + wf(s, t)|^{2} ds dt$$

$$= \mathcal{R}_{2} - \lim_{\boldsymbol{\varepsilon},\boldsymbol{\eta}\to\boldsymbol{0}} \frac{1}{16\pi^{2}\varepsilon\eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \{2 + we^{-i(\boldsymbol{u}x_{1}+\boldsymbol{v}x_{2})} + \overline{w}e^{i(\boldsymbol{u}x_{1}+\boldsymbol{v}x_{2})}\}$$

$$\cdot |\Delta_{\varepsilon,\eta}s(\boldsymbol{u}, \boldsymbol{v}; f)|^{2} du dv. \qquad (3.15)$$

 $\it Proof.$  It follows from Lemma 3, Theorem 3 and Minkowski's inequality that

$$\begin{split} \mathcal{R}_{1} \cdot \lim_{S,T \to \infty} \left\{ \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} |f(x_{1}+s, x_{2}+t) + wf(s, t)|^{2} ds dt \right\}^{1/2} \\ &= \mathcal{R}_{2} \cdot \lim_{\epsilon,\eta \to 0} \left\{ \frac{1}{16\pi^{2}\epsilon\eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |\Delta_{\epsilon,\eta} s_{x_{1},x_{2}}(u, v ; f) \right. \\ &+ w\Delta_{\epsilon,\eta} s(u, v ; f)|^{2} du dv \right\}^{1/2} \\ &\leq \mathcal{R}_{2} \cdot \lim_{\epsilon,\eta \to 0} \left\{ \frac{1}{16\pi^{2}\epsilon\eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |\Delta_{\epsilon,\eta} s_{x_{1},x_{2}}(u, v ; f) \right. \\ &- e^{i(ux_{1}+vx_{2})} \Delta_{\epsilon,\eta} s(u, v ; f)|^{2} du dv \right\}^{1/2} \\ &+ \mathcal{R}_{2} \cdot \lim_{\epsilon,\eta \to 0} \left[ \frac{1}{16\pi^{2}\epsilon\eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \{2 + we^{-i(ux_{1}+vx_{2})} + \overline{w}e^{i(ux_{1}+vx_{2})} \} \\ &\cdot |\Delta_{\epsilon,\eta} s(u, v ; f)|^{2} du dv \right]^{1/2}. \end{split}$$

Thus, by Lemma 5, (3.15) is proved.

Proof of THEOREM 4. By the Schwarz inequality and Lemma 1, we have

$$\begin{aligned} |\phi(x_1, x_2; f)| &\leq \mathcal{R}_1 \cdot \overline{\lim_{s, T \to \infty}} \left\{ \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} |f(x_1 + s, x_2 + t)|^2 ds dt \right\}^{1/2} \\ &\cdot \left\{ \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} |f(s, t)|^2 ds dt \right\}^{1/2} \\ &= \phi(0, 0; f) \,. \end{aligned}$$

Thus, (2.13) is proved.

Proof of THEOREM 5. By the Schwarz inequality, we have

$$\begin{split} |\phi(x_{1}+\varepsilon, x_{2}+\eta; f) - \phi(x_{1}, x_{2}; f)| \\ & \leq \mathcal{R}_{1} \cdot \lim_{S, T \to \infty} \left\{ \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} |f(x_{1}+\varepsilon+s, x_{2}+\eta+t) - f(x_{1}+s, x_{2}+t)|^{2} ds dt \right\}^{1/2} \\ & \cdot \left\{ \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} |f(s, t)|^{2} ds dt \right\}^{1/2}. \end{split}$$

If we now appeal to Lemmas 1, 3 and 4, then

$$\begin{split} \mathcal{R}_{1} - \lim_{S,T \to \infty} \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} |f(x_{1} + \varepsilon + s, x_{2} + \eta + t) - f(x_{1} + s, x_{2} + t)|^{2} ds dt \\ &= \mathcal{R}_{1} - \lim_{S,T \to \infty} \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} |f(x_{1} + \varepsilon + s, x_{2} + \eta + t)|^{2} ds dt \\ &+ \mathcal{R}_{1} - \lim_{S,T \to \infty} \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} |f(x_{1} + s, x_{2} + \eta + t)|^{2} ds dt \\ &- \mathcal{R}_{1} - \lim_{S,T \to \infty} \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} f(x_{1} + \varepsilon + s, x_{2} + \eta + t) \overline{f(x_{1} + s, x_{2} + t)} ds dt \end{split}$$

$$-\mathcal{R}_{1} \cdot \lim_{s, \tau \to \infty} \frac{1}{4ST} \int_{-\tau}^{T} \int_{-s}^{s} f(x_{1}+s, x_{2}+t) \overline{f(x_{1}+\varepsilon+s, x_{2}+\eta+t)} ds dt$$
  
=2\phi(0, 0; f)-\phi(\varepsilon, \eta; f)-\phi(-\varepsilon, -\eta; f).

Thus, we have

$$\begin{aligned} |\phi(x_1 + \varepsilon, x_2 + \eta; f) - \phi(x_1, x_2; f)| \\ &\leq [\phi(0, 0; f) \{2\phi(0, 0; f) - \phi(\varepsilon, \eta; f) - \phi(-\varepsilon, -\eta; f)\}]^{1/2}, \end{aligned}$$
(3.16)

from which the theorem follows immediately.

**Proof** of THEOREM 6. If we take w in (3.15) of Lemma 6 successively to equal  $\pm 1$ ,  $\pm i$ , and combine four expressions, then we obtain (2.14) by (3.2).

Finally, we prove Theorems 7 and 8.

**Proof** of THEOREM 7. Let  $f(x_1, x_2)$  belong to the class  $S(R^2)$ . Now, let us define

$$\phi_{\varepsilon,\eta}(x_1, x_2; f) = \frac{1}{16\pi^2 \varepsilon \eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{i(ux_1+vx_2)} |\Delta_{\varepsilon,\eta}s(u, v; f)|^2 du dv.$$
(3.17)

Then, it follows at once that

$$|\phi_{\epsilon,\eta}(x_1, x_2; f)| \leq \phi_{\epsilon,\eta}(0, 0; f).$$

Inasmuch as  $\phi_{\epsilon,\eta}(0,0;f)$  tends to  $\phi(0,0;f)$  as  $\varepsilon,\eta \rightarrow 0(\mathcal{R}_2)$ ,  $\phi_{\epsilon,\eta}(x_1,x_2;f)$  is bounded for all  $(x_1,x_2)\in \mathbb{R}^2$  and small values of  $\varepsilon,\eta$ . It, therefore, tends boundedly to  $\phi(x_1,x_2;f)$  as  $\varepsilon,\eta \rightarrow 0(\mathcal{R}_2)$ . Thus, by the bounded convergence theorem and Fubini's theorem, we get

$$\frac{1}{\lambda} \int_{-\lambda}^{\lambda} \left( 1 - \frac{|x_1|}{\lambda} \right) \phi(x_1, 0; f) dx_1$$
  
=  $\mathcal{R}_2 \cdot \lim_{\epsilon, \eta \to 0} \frac{1}{\lambda} \int_{-\lambda}^{\lambda} \left( 1 - \frac{|x_1|}{\lambda} \right) \phi_{\epsilon, \eta}(x_1, 0; f) dx_1$   
=  $\mathcal{R}_2 \cdot \lim_{\epsilon, \eta \to 0} \frac{1}{16\pi^2 \epsilon \eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{4 \sin^2 \frac{u\lambda}{2}}{u^2 \lambda^2} |\Delta_{\epsilon, \eta} s(u, v; f)|^2 du dv$ 

Hence, if  $f(x_1, x_2)$  belongs to the class  $S'(R^2)$ , then we have

$$\phi(0,0;f) = \lim_{\lambda \to 0} \frac{1}{\lambda} \int_{-\lambda}^{\lambda} \left(1 - \frac{|x_1|}{\lambda}\right) \phi(x_1,0;f) dx_1 ,$$

and

$$\lim_{\lambda \to 0} \mathcal{R}_{2} \lim_{\epsilon, \eta \to 0} \frac{1}{16\pi^{2} \epsilon \eta} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \left[ 1 - \frac{4 \sin^{2} \frac{u \lambda}{2}}{u^{2} \lambda^{2}} \right] |\Delta_{\epsilon, \eta} s(u, v; f)|^{2} du dv = 0.$$
(3.18)

Since, when  $|u\lambda| > \pi$ ,

$$1 - \frac{4\sin^2\frac{u\lambda}{2}}{u^2\lambda^2} > 1 - \frac{4}{\pi^2}$$

it follows from the positiveness of the integrand in (3.18) that

$$\lim_{\lambda \to 0} \mathcal{R}_{2^{-}} \overline{\lim_{\epsilon,\eta \to 0}} \frac{1}{16\pi^{2} \epsilon \eta} \int_{-\infty}^{\infty} \left[ \int_{\pi/\lambda}^{\infty} + \int_{-\infty}^{-\pi/\lambda} \right] |\Delta_{\epsilon,\eta} s(u,v;f)|^{2} du dv = 0$$

or simply

$$\lim_{A\to\infty} \mathcal{R}_2 \cdot \lim_{\epsilon,\eta\to 0} \frac{1}{16\pi^2 \varepsilon \eta} \int_{-\infty}^{\infty} \left[ \int_{A}^{\infty} + \int_{-\infty}^{-A} \right] |\Delta_{\epsilon,\eta} s(u,v;f)|^2 du dv = 0,$$

and we obtain (2.15). Similarly, we obtain (2.16).

Again, let  $f(x_1, x_2)$  belong to the class  $S(\mathbb{R}^2)$  and let (2.15) and (2.16) hold. Then, by (2.14) of Theorem 6,

$$\begin{split} |\phi(x_1, x_2; f) - \phi(0, 0; f)| \\ & \leq 2 \mathcal{R}_2 \cdot \overline{\lim_{\epsilon, \eta \to 0}} \frac{1}{16\pi^2 \epsilon \eta} \int_{-\infty}^{\infty} \left[ \int_A^{\infty} + \int_{-\infty}^{-A} \right] |\Delta_{\epsilon, \eta} S(u, v; f)|^2 du dv \\ & + 2 \mathcal{R}_2 \cdot \overline{\lim_{\epsilon, \eta \to 0}} \frac{1}{16\pi^2 \epsilon \eta} \left[ \int_A^{\infty} + \int_{-\infty}^{-A} \right] \int_{-\infty}^{\infty} |\Delta_{\epsilon, \eta} S(u, v; f)|^2 du dv \\ & + \mathcal{R}_2 \cdot \overline{\lim_{\epsilon, \eta \to 0}} \frac{1}{16\pi^2 \epsilon \eta} \int_{-A}^{A} \int_{-A}^{A} |e^{i(ux_1 + vx_2)} - 1| |\Delta_{\epsilon, \eta} S(u, v; f)|^2 du dv \\ & = I_1 + I_2 + I_3, \quad \text{say.} \end{split}$$

Now, let us choose A, as is possible by (2.15) and (2.16), so large that  $I_1$  and  $I_2$  do not exceed  $\delta/3$ . Let us then choose  $|x_1|$  and  $|x_2|$  so small that over  $(-A, A) \times (-A, A)$ 

$$|e^{i(ux_1+vx_2)}\!-\!1|\!\leq\!rac{\delta}{3\phi(0,0\,;\,f)}$$
 ,

and, therefore,  $I_3$  does not exceed  $\delta/3$ . Thus, since we have

$$|\phi(x_1, x_2; f) - \phi(0, 0; f)| \leq \delta$$
,

it follows from Theorem 5 that  $f(x_1, x_2)$  belongs to the class  $S'(R^2)$ . This completes the proof of Theorem 7.

**Proof** of THEOREM 8. From Theorem 4 and Definition 3, it follows immediately that  $\phi(x_1, x_2; f)$  is bounded and continuous on  $\mathbb{R}^2$ . Also,

$$\overline{\phi(-x_1, -x_2; f)} = \phi(x_1, x_2; f)$$

for all  $(x_1, x_2) \in \mathbb{R}^2$ . Furthermore, for any finite sequence  $\{(\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n)\}$  of points in  $\mathbb{R}^2$  and any finite sequence  $\{z_1, \dots, z_n\}$  of complex numbers,

$$\begin{split} \sum_{\mu,\nu=1}^{n} \phi(\alpha_{\mu} - \alpha_{\nu}, \beta_{\mu} - \beta_{\nu}) z_{\mu} \overline{z}_{\nu} \\ &= \mathcal{R}_{1} \cdot \lim_{S,T \to \infty} \frac{1}{4ST} \sum_{\mu,\nu=1}^{n} \int_{-T}^{T} \int_{-S}^{S} f(\alpha_{\mu} - \alpha_{\nu} + s, \beta_{\mu} - \beta_{\nu} + t) \overline{f(s,t)} ds dt z_{\mu} \overline{z}_{\nu} \\ &= \mathcal{R}_{1} \cdot \lim_{S,T \to \infty} \frac{1}{4ST} \int_{-T}^{T} \int_{-S}^{S} \left| \sum_{\mu=1}^{n} f(\alpha_{\mu} + s, \beta_{\mu} + t) z_{\mu} \right|^{2} ds dt \ge 0 \,. \end{split}$$

Consequently,  $\phi(x_1, x_2; f)$  is positive definite. Thus, by Bochner's representation theorem and Lévy's inversion formula, there exists a measure  $\mu$  on  $(R^2, \mathcal{B}_2)$  such that  $\mu(R^2) = \phi(0, 0; f)$ , and (2.17) and (2.18) hold with  $\phi(x_1, x_2; f)$  instead of  $\phi(x_1, x_2)$ .

Now, let us put

$$A(u, v) = \int_{-\infty}^{v} \int_{-\infty}^{u} d\mu(\xi_1, \xi_2) .$$
 (3.19)

Then,  $\Lambda(u, v)$  is monotone increasing and

$$\int_{-\infty}^{\infty}\int_{-\infty}^{\infty}|dA(u,v)| = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty}d\mu(\xi_1,\xi_2) = \mu(R^2) = \phi(0,0;f).$$

Moreover, by (2.17) and (2.18) with  $\phi(x_1, x_2; f)$  instead of  $\phi(x_1, x_2)$ , we have (2.20) and (2.21). This completes the proof.

Remark. The generalized harmonic analysis we established in this paper was based on the two-dimensional Wiener formula (Theorem 2 or 2'). Its limit process, therefore, also depended on the limit process of the above two-dimensional Wiener formula. On the other hand, the two-dimensional Wiener formula is also proved under the unrestricted rectangular mean concerning the double limit process. For details, refer to Pitt [8] and Rudin [9]. Thus, if we base the argument on this sort of two-dimensional Wiener formula, then the generalized harmonic analysis also holds under the above limit process instead of the restricted limit process.

### Acknowledgement

It is a pleasure to acknowledge Professor Sumiyuki Koizumi of Keio University, who suggested me the problem treated here, and was consistently helpful in the course of preparation of the present paper.

#### REFERENCES

- [1] ANZAI, K., KOIZUMI, S. and MATSUOKA, K. (to appear): On the Wiener formula of functions of two variables, *Tokyo Jour. Math.*
- [2] BOCHNER, S. (1959): Lectures on Fourier integrals, Ann. of Math. Studies, No. 42, Princeton Univ. Press, New Jersey.
- [3] KAWATA, T. (1972): Fourier analysis in probability theory, Academic Press, New York.
- [4] KOIZUMI, S. (1959): On the singular integrals VI, Proc. Japan Acad., 35, 323-328.

- [5] KOIZUMI, S. (1959): On the Hilbert transform I, Jour. Faculty Sci. Hokkaido Univ., Series I, 14, 153-224.
- [6] KOIZUMI, S. (1962): Hilbert transform in the Stepanoff space, Proc. Japan Acad., 38, 735-740.
- [7] MATSUOKA, K. (to appear): On the generalized Hilbert transforms of functions of two variables.
- [8] PITT, H. R. (1938): General Tauberian theorems, Proc. London Math. Soc., 44, 243-288.
- [9] RUDIN, W. (1960): Fourier analysis on groups, Interscience Tracts in Pure and Applied Math., No. 12, New York.
- [10] WIENER, N. (1930): Generalized harmonic analysis, Acta Math., 55, 117-258.
- [11] WIENER, N. (1933): The Fourier integral and certain of its applications, Cambridge Univ. Press, London.