

Title	Enzymatic formation of 5-fluorouracil from 1-(tetrahydro-2-furanyl)-5-fluorouracil (tegafur) in human tumor tissues
Sub Title	
Author	河野, 彬(Kono, Akira) 原, 泰寛(Hara, Yasuhiro) 松島, 美一(Matsushima, Yoshikazu)
Publisher	共立薬科大学
Publication year	1982
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.27 (1982.) ,p.78- 78
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000027-0078

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Enzymatic Formation of 5-Fluorouracil from 1-(Tetrahydro-2-furanyl)-5-fluorouracil (Tegafur) in Human Tumor Tissues*

AKIRA KONO,** YASUHIRO HARA,** and YOSHIKAZU MATSUSHIMA

河野 彬,** 原 泰寛,** 松島美一

1-(Tetrahydro-2-furanyl)-5-fluorouracil (Tegafur) has an antitumor activity with a wide spectrum of activity similar to 5-fluorouracil (FU) and is considered to be a chemical depot form of FU. It has been believed that FU is generated *in vivo* from Tegafur by hepatic metabolism involving cytochrome P-450.

We found that a soluble fraction of human lung cancer catalyzed the phosphorolytic cleavage of Tegafur. The catalysis was suppressed in the presence of excess thymidine, but not in the presence of 1-(2'-deoxy- β -D-glucopyranosyl)-thymine, an inhibitor of uridine phosphorylases.

The phosphorolytic activities of Tegafur and thymidine were much higher in extracts of tumor tissues than in those of normal tissues of human lung. The Tegafur phosphorolytic activity was hardly present in extract of mouse tumor (Sarcoma 180), which showed low thymidine phosphorylase activity.

We assume that the cleavage of Tegafur to FU is catalyzed by a thymidine phosphorylase activity and represents a possible activation mechanism of Tegafur in human tumor tissues.

* 本報告は *Chem. Pharm. Bull.*, 29, 1486~1488 (1981) に発表.

** 九州がんセンター.