

Title	A reticuloendothelial system-activating arabinoxylan from the bark of cinnamomum cassia
Sub Title	
Author	金成, 美枝子(Kanari, Mieko) 友田, 正司(Tomoda, Masashi) 権田, 良子(Gonda, Ryoko) 清水, 訓子(Shimizu, Noriko) 木村, 都(Kimura, Miyako) 川口, 雅美(Kawaguchi, Masami) 川辺, 千織(Kawabe, Chiori)
Publisher	共立薬科大学
Publication year	1989
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.34 (1989. ) ,p.61- 61
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	<a href="https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000034-0061">https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000034-0061</a>

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

## A Reticuloendothelial System-Activating Arabinoxylan from the Bark of *Cinnamomum cassia*\*

Mieko KANARI, Masashi TOMODA, Ryōko GONDA, Noriko SHIMIZU,  
Miyako KIMURA, Masami KAWAGUCHI and Chiori KAWABE

金成美枝子, 友田正司, 権田良子, 清水訓子  
木村 都, 川口雅美, 川辺千織

The dried bark of *Cinnamomum cassia* is a very famous crude drug, Cinnamomi Cortex. We have isolated a neutral polysaccharide, named cinnaman AX, having high immunological activity on the RES from the hot water extract of this crude drug by treatment with sodium lauryl sulfate, sodium chloride and acetone followed by gel chromatography on Toyopearl HW-65F, Sephacryl S-500 and Cellulofine GCL-25 m columns. The polysaccharide gave a single spot on glass-fiber paper electrophoresis and gave a single peak on gel chromatography.

Cinnaman AX is composed of L-arabinose: D-xylose in the molar ratio of 4:3, and its molecular mass was estimated to be about  $1.0 \times 10^6$ .

The results of methylation analysis, periodate oxidation and  $^{13}\text{C}$ -NMR spectrum suggested that the minimal unit of cinnaman AX is composed of twenty terminal  $\alpha$ -L-arabinofuranose, one terminal  $\beta$ -L-arabinopyranose, twenty  $\beta$ -1,3-linked L-arabinopyranose, ten  $\beta$ -1,4-linked D-xylopyranose and twenty-one 3,4-branched  $\beta$ -D-xylopyranose units.

In addition, the controlled Smith degradation and partial hydrolysis studies revealed that cinnaman AX has a backbone chain composed of  $\beta$ -1,4-linked D-xylopyranose units. About 70% of the xylose residues in the backbone carry side chains composed of  $\alpha$ -L-Araf-(1 $\rightarrow$ 3)- $\beta$ -L-Araf and  $\alpha$ -L-Araf-(1 $\rightarrow$ 4)- $\beta$ -D-Xylp-(1 $\rightarrow$ 3)- $\beta$ -L-Araf units at position 3. There is one branching unit having a terminal  $\beta$ -L-arabinopyranose per twenty or twenty-six branching residues bearing the disaccharide and trisaccharide side chains.

The effect of cinnaman AX on a RES was demonstrated by the *in vivo* carbon clearance test. When administered i. p. (5, 20 and 50 mg/kg), the phagocytic indices of cinnaman AX were  $0.1310 \pm 0.0117$ ,  $0.1877 \pm 0.0443$  and  $0.3120 \pm 0.1054$ . Thus the values were remarkably increased, suggesting powerful activation of RES. Cinnaman AX is a new structural type of polysaccharide having a remarkable activity on the RES. The prominent presence of  $\beta$ -L-arabinopyranose units is especially characteristic.

---

\* 本報告は *Chem. Pharm. Bull.*, 37, 3191—3194 (1989) に発表.