

Title	An acidic polysaccharide having activity on the reticuloendothelial system from the roots and rhizomes of <i>saposhnikovia divaricata</i>
Sub Title	
Author	清水, 訓子(Shimizu, Noriko) 友田, 正司(Tomododa, Masashi) 権田, 良子(Gonda, Ryoko) 金成, 美枝子(Kanari, Mieko) 久保田, 明子(Kubota, Akiko) 久保田, 朱美(Kubota, Akemi)
Publisher	共立薬科大学
Publication year	1989
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.34 (1989.) ,p.62- 62
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000034-0062

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**An Acidic Polysaccharide Having Activity on the Reticuloendothelial System
from the Roots and Rhizomes of *Saposhnikovia divaricata****

Noriko SHIMIZU, Masashi TOMODA, Ryōko GONDA, Mieko KANARI,
Akiko KUBOTA and Akemi KUBOTA

清水訓子, 友田正司, 権田良子, 金成美枝子
久保田明子, 久保田朱美

From the hot water extract of the roots and rhizomes of *Saposhnikovia divaricata*, an acidic polysaccharide, named saposhnikovan C, was isolated by fractionation with cetyltrimethylammonium bromide and on a Sephadex G-50 column followed by chromatography on DEAE-Sephadex A-25 (acetate) and on Sephacryl S-300 columns. The polysaccharide gave a single spot on cellulose acetate membrane electrophoresis and gave a single band on PAGE. In addition, it gave a single peak on gel chromatography.

Saposhnikovan C is composed of L-arabinose: D-galactose: L-rhamnose: D-galacturonic acid: acetyl: methoxyl groups in the molar ratio of 8:8:7:27:7:8, and the molecular mass was estimated to be 132000. About 30% of the D-galacturonic acid residues exist as the methyl esters.

The results of methylation analysis of the original polysaccharide and the carboxyl-reduced derivative and ^{13}C -NMR spectrum suggested that the minimal unit of saposhnikovan C is composed of four terminal α -L-arabinofuranose, two α -1,5-linked L-arabinofuranose, two 3,5-branched α -L-arabinofuranose, five terminal β -D-galactopyranose, one β -1,4-linked D-galactopyranose, two 3,4-branched β -D-galactopyranose, four α -1,2-linked L-rhamnopyranose, three 2,4-branched α -L-rhamnopyranose, two 3,4-branched α -D-galactopyranosyluronic acid and twenty-five α -1,4-linked D-galacturonan units. In addition, the controlled Smith degradation and partial hydrolysis studies revealed the presence of a backbone chain consisting of α -1,4-linked D-galacturonic acid residues with interspersed α -1,2-linked L-rhamnose residues and of side chains composed of 1,4-linked and 3,4-branched β -D-galactan and 3,5-branched α -L-arabinan.

The effect of saposhnikovan C on a RES was demonstrated by the *in vivo* carbon clearance test. When administered i. p. (50 mg/kg), the phagocytic index was 0.3244 ± 0.0662 . Thus it showed remarkable RES-potentiating activity. It is interesting that saposhnikovan C having highly immunological activity possesses a pectin-like rhamnogalacturonan backbone, while structurally typical pectins show no RES activity.

* 本報告は *Chem. Pharm. Bull.*, 37, 3054—3057 (1989) に発表.