

| | |
|------------------|---|
| Title | Characterization of polysaccharides having activity on the reticuloendothelial system from the rhizome of curcuma longa |
| Sub Title | |
| Author | 権田, 良子(Gonda, Ryoko) 友田, 正司(Tomododa, Masashi) 清水, 訓子(Shimizu, Noriko) 金成, 美枝子(Kanari, Mieko) |
| Publisher | 共立薬科大学 |
| Publication year | 1990 |
| Jtitle | 共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.35 (1990.) ,p.34- 34 |
| JaLC DOI | |
| Abstract | |
| Notes | 抄録 |
| Genre | Technical Report |
| URL | https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0034 |

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Characterization of Polysaccharides Having Activity
on the Reticuloendothelial System from
the Rhizome of *Curcuma longa****

Ryōko GONDA, Masashi TOMODA, Noriko SHIMIZU and Mieko KANARI

権田良子, 友田正司, 清水訓子, 金成美枝子

The crude polysaccharide fraction was isolated from the rhizome of *Curcuma longa* by hot water extraction followed by precipitation with ethanol. It was fractionated by DEAE-Sephadex A-25 (carbonate) column chromatography, and the fraction having high activity was subjected to affinity chromatography on Con A-Sepharose and gel chromatography on Sephacryl S-300. Three polysaccharides, named ukonan A, ukonan B and ukonan C were obtained. They were homogeneous on electrophoresis and gel chromatography, and showed remarkable reticuloendothelial system (RES)-potentiating activity in a carbon clearance test even in a minute dose (5 mg per kg).

Among them, ukonan B is the major one in terms of the activity. It contains L-arabinose, D-xylose, D-galactose, D-glucose, L-rhamnose and D-galacturonic acid in the molar ratio of 12:4:12:1:2:4 in addition to 6.6% peptide moiety. The value of 3.3×10^4 was estimated for its molecular mass.

Methylation analysis and ^{13}C -NMR studies indicated that the minimal unit of ukonan B is composed of twenty-four terminal α -L-arabinofuranose, ten α -1,3-linked L-arabinopyranose, two α -1,5-linked L-arabinofuranose, two β -1,4-linked D-xylose, ten β -3,4-branched D-xylose, four terminal β -D-galactose, ten β -1,3-linked D-galactose, four β -1,6-linked D-galactose, eighteen β -3,6-branched D-galactose, two terminal α -D-glucose, one α -1,4-linked D-glucose, four α -1,2-linked L-rhamnose, two α -2,4-branched L-rhamnose and twelve α -1,4-linked D-galacturonic acid residues. Thus, ukonan B may be basically a kind of α -L-arabino- β -3,6-branched D-galactan.

* 本報告は *Chem. Pharm. Bull.*, 38, 482-486 (1990) に発表.