

Title	A novel neutral polysaccharide having activity on the reticuloendothelial system from the root of glycyrrhiza uralensis
Sub Title	
Author	清水, 訓子(Shimizu, Noriko) 友田, 正司(Tomoda, Masashi) 金成, 美枝子(Kanari, Mieko) 権田, 良子(Gonda, Ryoko) 佐藤, 明美(Sato, Akemi) 佐藤, 希子(Sato, Noriko)
Publisher	共立薬科大学
Publication year	1990
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.35 (1990.) ,p.43- 43
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0043

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**A Novel Neutral Polysaccharide Having Activity on
the Reticuloendothelial System from the
Root of *Glycyrrhiza uralensis****

Noriko SHIMIZU, Masashi TOMODA, Mieko KANARI, Ryōko GONDA,
Akemi SATOH and Noriko SATOH

清水訓子, 友田正司, 金成美枝子, 権田良子, 佐藤明美, 佐藤希子

A novel neutral polysaccharide; named glycyrrhizan UC, was isolated from the root of *Glycyrrhiza uralensis* by hot water extraction followed by treatment with cetyltrimethylammonium bromide, gel chromatography with Sephacryl S-300, and affinity chromatography on Con A-Sepharose. The polysaccharide was obtained from the fraction retained in Con A. It produced a single band on PAGE, and a single peak on gel chromatography. A value of 6.9×10^4 was yielded for the molecular mass.

Glycyrrhizan UC is composed of L-arabinose : D-galactose : D-glucose : L-rhamnose in the molar ratio of 10:30:27:1. Its effect on the reticuloendothelial system was demonstrated by a carbon clearance test. The phagocytic index was significantly increased, and the value was superior than those of glycyrrhizans UA and UB.

Chemical and spectroscopic studies indicated that the minimal unit of glycyrrhizan UC is composed of seven terminal α -L-arabinofuranose, seven α -1,5-linked L-arabinofuranose, two terminal α -L-arabinopyranose, four α -1,3-linked L-arabinopyranose, four terminal β -D-galactopyranose, ten β -1,3-linked D-galactopyranose, eighteen β -1,4-linked D-galactopyranose, ten β -1,6-linked D-galactopyranose, twelve β -3,6-branched D-galactopyranose, five β -4,6-branched D-galactopyranose, one β -2,4-branched D-galactopyranose, twelve terminal α -D-glucopyranose, eight α -1,3-linked D-glucopyranose, twenty-six α -1,4-linked D-glucopyranose, one α -3,6-branched D-glucopyranose, seven α -4,6-branched D-glucopyranose, one terminal α -L-rhamnopyranose and one α -1,2-linked L-rhamnopyranose residues.

* 本報告は *Chem. Pharm. Bull.*, 38, 3069—3071 (1990) に発表.