Keio Associated Repository of Academic resouces

Reto / Baselated Repositor y of / Reddefine resources	
Title	Effects of synthetic estrogens, (+)-, (-)-, dl- and meso-hexestrol stereoisomers on microtubule assembly
Sub Title	
Author	榊原, 由美子(Sakakibara, Yumiko) 長谷川, 清(Hasegawa, Kiyoshi) 小田, 泰子(Oda, Taiko) 斎藤, 肇(Saito, Hajime) 児玉, 昌彦(Kodama, Masahiko) 平田, 愛子(Hirata, Aiko) 松橋, 通生(Matsuhashi, Michio) 佐藤, 良博(Sato, Yoshihiro)
Publisher	共立薬科大学
Publication year	1990
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.35 (1990.), p.57-57
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000035-0057

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって 保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Effects of Synthetic Estrogens, (+)-, (-)-, dl- and meso-Hexestrol Stereoisomers on Microtubule Assembly

Yumiko Sakakibara, Kiyoshi Hasegawa, Taiko Oda, Hazime Saitô**, Masahiko Kodama**, Aiko Hirata***, Michio Matsuhashi*** and Yoshihiro Sato

榊原由美子, 長谷川清, 小田泰子, 斎藤 肇**, 児玉昌彦**, 平田愛子***, 松橋通生***, 佐藤良博

We previously reported on the inhibition of microtubule polymerization and the formation of ribbon structures by synthetic estrogens. The present investigation aimed to analyze these effects in vitro on stereochemical point of view, using hexestrol isomers ((+)-hexestrol, (-)-hexestrol and meso-hexestrol) and dl-hexestrol. Among hexestrols, dl-hexestrol showed the highest activity in ribbon formation from microtubule proteins at $100\,\mu\text{M}$.

On the other hand, meso-hexestrol was distinguished from others by inhibition of microtubule assembly and formation of a large amount of aggregates from purified tubulin in the presence of $MgCl_2$ and dimethyl sulfoxide. These results were discussed with physico-chemical properties of hexestrols, e.g. absolute configurations as well as circular dichroism spectra and solid state carbon-13 nuclear magnetic resonance spectra.

^{*} 本報告は Biochem. Phamacol., 39, 167-172 (1990) に発表.

^{**} 国立がんセンター研究所生物物理部

^{***} 東大応微研第一研究部