

Title	Characterization of a polysaccharide having activity on the reticuloendothelial system from the stolon of <i>glycyrrhiza glabra</i> var. <i>glandulifera</i>
Sub Title	
Author	清水, 訓子(Shimizu, Noriko) 友田, 正司(Tomoda, Masashi) 佐藤, 麻希子(Sato, Makiko) 権田, 良子(Gonda, Ryoko) 大原, 直子(Ohara, Naoko)
Publisher	共立薬科大学
Publication year	1991
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.36 (1991.) ,p.41- 41
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000036-0041

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Characterization of a Polysaccharide Having Activity on
the Reticuloendothelial System from the Stolon
of *Glycyrrhiza glabra* var. *glandulifera****

Noriko SHIMIZU, Masashi TOMODA, Makiko SATOH, Ryōko GONDA
and Naoko ŌHARA

清水訓子, 友田正司, 佐藤麻希子, 権田良子, 大原直子

An acidic polysaccharide, named glycyrrhizan GA, was isolated from the stolon of *Glycyrrhiza glabra* var. *glandulifera*. It produced a single band on electrophoresis and a single peak on gel chromatography, and its molecular mass was estimated to be 85,000.

Glycyrrhizan GA is composed of L-arabinose : D-galactose : L-rhamnose : D-galacturonic acid : D-glucuronic acid in the molar ratio of 22 : 10 : 1 : 2 : 1, in addition to small amounts of O-acetyl groups. Part of the hexuronic acid residues exist as methyl esters.

Chemical and spectroscopic studies indicated that the minimal unit of polysaccharide is composed of thirteen terminal α -L-arabinofuranose, four α -1,3-linked L-arabinofuranose, twenty-two α -1,5-linked L-arabinofuranose, five α -2,5-branched L-arabinofuranose, one α -1,2-linked L-rhamnopyranose, one α -2,4-branched L-rhamnopyranose, one terminal β -D-galactopyranose, six β -1,3-linked D-galactopyranose, three β -1,6-linked D-galactopyranose, ten β -3,6-branched D-galactopyranose, four α -1,4-linked D-galactopyranosyluronic acid and terminal β -D-glucopyranosyluronic acid residues.

Glycyrrhizan GA showed remarkable reticuloendothelial system-potentiating activity in a carbon clearance test.

* 本報告は *Chem. Pharm. Bull.*, 39, 2082—2086 (1991) に発表。