

Title	Effect of meso-hexestrol, a synthetic estrogen, on S-tubulin
Sub Title	
Author	榊原, 由美子(Sakakibara, Yumiko) 小田, 泰子(Oda, Taiko) 平田, 愛子(Hirata, Aiko) 松橋, 通生(Matsuhashi, Michio) 佐藤, 良博(Sato, Yoshihiro)
Publisher	共立薬科大学
Publication year	1991
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.36 (1991. ) ,p.53- 53
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	<a href="https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000036-0053">https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000036-0053</a>

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Effect of meso-Hexestrol, a Synthetic Estrogen, on S-Tubulin\***

Yumiko SAKAKIBARA, Taiko ODA, Aiko HIRATA\*\*, Michio MATSUHASHI\*\*,  
and Yoshihiro SATO

榑原由美子, 小田泰子, 平田愛子\*\*, 松橋通生\*\*, 佐藤良博

We have reported that meso-hexestrol, a synthetic estrogen, inhibits microtubule assembly and induces microtubule proteins into twisted ribbon structures. On the other hand, Serrano *et al.* proved that S-tubulin, which lacks the C-terminal moiety of tubulin subunits, assembles into sheet structures in the absence of microtubule-associated proteins (MAPs). In the present investigation, we attempted to clarify whether meso-hexestrol could induce the ribbon structure from S-tubulin. meso-Hexestrol delayed the initiation of polymerization of S-tubulin into sheet structures in a dose-dependent manner below 50  $\mu$ M.

But the effect of meso-hexestrol on S-tubulin was reduced in the presence of either tau or microtubule-associated protein 2 (MAP2) in a MAPs-concentration-dependent manner. At concentrations higher than 100  $\mu$ M, meso-hexestrol inhibited the polymerization of S-tubulin into sheet structures, without forming ribbon structures. The present results may indicate that meso-hexestrol interacts with S-tubulin, and its interaction is affected by MAPs.

\* 本報告は *Chem. Pharm. Bull.*, 38, 3419—3422 (1990) に発表.

\*\* 東大応微研第一研究部