

Title	Lysophospholipase L ₁ from escherichia coli K-12 overproducer
Sub Title	
Author	唐沢, 健(Karasawa, Ken) 工藤, 一郎(Kudo, Ichiro) 小林, 哲幸(Kobayashi, Tetsuyuki) 本間, 浩(Honma, Hiroshi) 知場, 伸介(Chiba, Nobuyoshi) 水島, 洋(Mizushima, Hiroshi) 井上, 圭三(Inoue, Keizo) 野島, 庄七(Nojima, Shoshichi)
Publisher	共立薬科大学
Publication year	1991
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.36 (1991.) ,p.60- 60
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000036-0060

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Lysophospholipase L₁ from *Escherichia coli* K-12 Overproducer

Ken Karasawa***, Ichiro Kudo**, Tetsuyuki Kobayashi*****, Hiroshi Homma,
Nobuyoshi Chiba**, Hiroshi Mizushima****, Keizo Inoue**,
and Shoshichi Nojima***

唐沢 健***, 工藤一郎**, 小林哲幸*****, 本間 浩, 知場伸介**,
水島 洋****, 井上圭三**, 野島庄七***

After screening 900 *E. coli* strains of the Clarke and Carbon collection for by lysophospholipase L₁ activities, we isolated a clone bearing the plasmid pLC6-34, which showed an increased level of lysophospholipase L₁ activity. Strains bearing the plasmid pC124, a subclone of pLC6-34 in plasmid vector pUC8, showed approximately 11.4 times higher lysophospholipase L₁ activity than that of the parental strain. Starting from those overproducing strains, the lysophospholipase L₁ was purified to near homogeneity by sequential use of ammonium sulfate fractionation, Sephacryl S-300, DEAE-cellulose, hydroxyapatite and Sephacryl S-200 column chromatographies. The apparent molecular weight of the purified lysophospholipase L₁ was estimated to be 20,500-22,000 both by SDS-polyacrylamide gel electrophoresis and by gel permeation chromatography. The specific activity of the homogeneous lysophospholipase L₁ was 10,400 nmol/min/mg protein when 1-acyl-sn-glycero-3-phosphoethanolamine was used as the substrate. The amino acid sequence of the amino-terminal portion of purified lysophospholipase L₁ was determined and was different from that of lysophospholipase L₂, which had previously been purified from the envelope fraction of *E. coli* strains bearing its cloned structural gene, *pldB* [Karasawa, K., Kudo, I., Kobayashi, T., Sa-eki, T., Inoue, K., & Nojima, S. (1985) *J. Biochem* 98, 1117—1125]. The gene responsible for overproduction of lysophospholipase L₁ activity was designated as *pldC* (*phospholipid degradation C*). Its restriction enzyme map was also different from that of cloned *pldB*. These results further confirmed that, in *E. coli*, there are two lysophospholipases with distinct characteristics.

* 本報告は *J. Biochem.*, **109**, 288—293 (1991) に発表.

** 東京大学薬学部

*** 帝京大学薬学部

**** 国立ガンセンター研究所生物物理部

***** 名古屋市立大学薬学部