

Title	Constituents of the seed of malva verticillata : VIII. Smith degradation of MVS-VI, the major acidic polysaccharide, and anti-complementary activity of products
Sub Title	
Author	友田, 正司(Tomoda, Masashi) 浅原, 初木(Asahara, Hatsuki) 権田, 良子(Gonda, Ryoko) 高田, 勝利(Takada, Katsutoshi)
Publisher	共立薬科大学
Publication year	1992
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.37 (1992.) ,p.54- 54
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000037-0054

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**Constituents of the Seed of *Malva verticillata*. VIII. Smith
Degradation of MVS-VI, the Major Acidic Polysaccharide,
and Anti-Complementary Activity of Products***

Masashi TOMODA, Hatsuki ASAHARA, Ryōko GONDA and Katsutoshi TAKADA

友田正司, 浅原初木, 権田良子, 高田勝利

The controlled Smith degradation of MVS-VI, the major acidic polysaccharide having remarkable anti-complementary activity isolated from the seeds of *Malva verticillata* L., was performed. Methylation analysis of both the primary and the secondary Smith degradation products indicated that the core structural features of MVS-VI include a backbone chain composed of β -1,3-linked D-galactose residues. The majority of galactose units in the backbone carry side chains composed of β -1,3 - and β -1,6-linked D-galactosyl residues at position 6. The controlled Smith degradation of MVS-VI afforded three products, and they showed considerable anti-complementary activity. This fact suggests that the core structure contributes to the activity. It is interesting that two products showed almost the same value on the activity despite a marked difference in molecular mass between them.

* 本報告は *Chem. Pharm. Bull.*, **40**, 2219—2221 (1992) に発表.