

Title	An acidic polysaccharide having immunological activities from the rhizome of cnidium officinale
Sub Title	
Author	友田, 正司(Tomoda, Masashi) 大原, 直子(Ohara, Naoko) 権田, 良子(Gonda, Ryoko) 清水, 訓子(Shimizu, Noriko) 高田, 勝利(Takada, Katsutoshi) 佐藤, 泰代(Sato, Yasuyo) 白井, 小百合(Shirai, Sayuri)
Publisher	共立薬科大学
Publication year	1992
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.37 (1992.) ,p.57- 57
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000037-0057

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the Keio Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

An Acidic Polysaccharide Having Immunological Activities from the Rhizome of *Cnidium officinale**

Masashi TOMODA, Naoko ŌHARA, Ryōko GONDA, Noriko SHIMIZU,
Katsutoshi TAKADA, Yasuyo SATOH and Sayuri SHIRAI

友田正司, 大原直子, 権田良子, 清水訓子,
高田勝利, 佐藤泰代, 白井小百合

An acidic polysaccharide, designated as cnidirhan AG, was isolated from the rhizomes of *Cnidium officinale* MAKINO. It was homogeneous on electrophoresis and gel chromatography, and its molecular mass was estimated to be 5.1×10^4 . It showed pronounced reticuloendothelial system-potentiating activity in a carbon clearance test, and had a remarkable effect on both anti-complementary and alkaline phosphatase-inducing activities. It is composed of L-arabinose : D-galactose : D-glucuronic acid in the molar ratio of 2 : 6 : 1, in addition to small amounts of O-acetyl groups. Methylation analysis, carbon-13 nuclear magnetic resonance, controlled Smith degradation and limited acid hydrolysis indicated that the core structural features of cnidirhan AG include a backbone chain composed of β -1,3-linked D-galactose residues. Some of the galactose units in the backbone carry β -D-galactosyl side chains at position 6. Both α -L-arabinosyl arabinose side chains and terminal β -D-glucuronic acid residues are linked to the core galactan units.

Cnidirhan AG can be classified under a novel structural type of acidic arabino-3,6-galactan having high immunological activity.

* 本報告は *Chem. Pharm. Bull.*, 40, 3025–3029 (1992) に発表.