

Title	pH-sensitive docosanoic acid-nicotinamide complex, and its applicability to a pH-responsive drug delivery system
Sub Title	
Author	横山, 祥子(Yokoyama, Shoko) 春原, 政明(Sunohara, Masaaki) 藤江, 忠雄(Fujie, Tadao)
Publisher	共立薬科大学
Publication year	1993
Jtitle	共立薬科大学研究年報 (The annual report of the Kyoritsu College of Pharmacy). No.38 (1993. ) ,p.85- 85
JaLC DOI	
Abstract	
Notes	抄録
Genre	Technical Report
URL	<a href="https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000038-0085">https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=AN00062898-00000038-0085</a>

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

**pH-Sensitive Docosanoic Acid–nicotinamide Complex, and Its  
Applicability to a pH-Responsive Drug Delivery System\***

Shoko YOKOYAMA, Masaaki SUNOHARA and Tadao FUJIE

横山祥子, 春原政明, 藤江忠雄

The release of nicotinamide (NAA) from docosanoic acid (C22)–NAA complex was measured in pH 1.2 — 6.8 aqueous media. The release of NAA was ON-state below pH 6.2 and OFF-state at pH 6.8, suggesting that the release behavior is applicable to a pH-responsive drug delivery system.

---

\* 本報告は *Chem. Pharm. Bull.*, 41, 1674—1676 (1993) に発表。