

Title	Utilization of sperm whale oil (preparation of oleyl oleate from sperm blubber oil and its hydroxy-derivatives)
Sub Title	
Author	木村, 勝紀(Kimura, Masanori)
Publisher	慶應義塾大学藤原記念工学部
Publication year	1966
Jtitle	Proceedings of the Fujihara Memorial Faculty of Engineering Keio University (慶應義塾大学藤原記念工学部研究報告). Vol.19, No.76 (1966.) ,p.268(50)- 268(50)
JaLC DOI	
Abstract	
Notes	Summaries of Doctor and Master Theses Master of Engineering, 1966
Genre	Departmental Bulletin Paper
URL	https://koara.lib.keio.ac.jp/xoonips/modules/xoonips/detail.php?koara_id=KO50001004-00190076-0050

慶應義塾大学学術情報リポジトリ(KOARA)に掲載されているコンテンツの著作権は、それぞれの著作者、学会または出版社/発行者に帰属し、その権利は著作権法によって保護されています。引用にあたっては、著作権法を遵守してご利用ください。

The copyrights of content available on the KeiO Associated Repository of Academic resources (KOARA) belong to the respective authors, academic societies, or publishers/issuers, and these rights are protected by the Japanese Copyright Act. When quoting the content, please follow the Japanese copyright act.

Utilization of Sperm Whale Oil

(Preparation of Oleyl Oleate from Sperm Blubber Oil and its Hydroxy-Derivatives)

Masanori KIMURA*

The crude oleyl oleate was prepared from sperm blubber oil by the following process.

The sperm blubber oil was lightly polymerized to remove the trace of higher unsaturated components, followed by the recrystallization and molecular distillation, then the crude oleyl oleate was obtained. Its characteristics were as follows; I. V. 80.9, Sap. V. 107.7.

Further, the purity of the prepared ester was checked by the gas-liquid chromatography and was compared with the synthesized oleyl oleate by the Swern's method (1). It was confirmed that the given ester was almost pure oleyl oleate, which contained ca. 10% of oleyl palmitate and cetyl oleate.

To find the new uses of oleyl oleate, the ester was hydroxylated with hydrogen peroxide. The constants of hydroxylated ester were as follows; m. p. 31—34°, I. V. 3.7, Sap. V. 142.8 and OH-V. 83.5

It was a mixture of hydroxy ester and a unknown ester occurred by the breakdown of the C-C double bonds.

*木 村 勝 紀